



Striped Cucumber Beetle

Pest Fact Sheet 20

Dr. Alan T. Eaton, Extension Specialist, Entomology

UNH Cooperative Extension Programs

	Community and Economic Development
	Food and Agriculture ✓
	Natural Resources
	Youth and Family

Introduction

The striped cucumber beetle, *Acalymma vittatum*, is one of the most devastating pests of cucurbits (cucumbers, summer and winter squashes, all types of melons and pumpkins) east of the Rocky Mountains. Both adults and larvae feed on cucurbit crops. This insect is also responsible for the spread of plant diseases such as bacterial wilt and squash mosaic virus. Cantaloupes and muskmelons are especially vulnerable to bacterial wilt spread by the beetles.

Description

The adult beetles are ¼" long and yellow-green with black longitudinal stripes. Eggs are small and orange-to-yellow in color. The worm-like larvae are slender, white, and about ½" long when full-grown.

Life Cycle

The insects overwinter as unmated adults in the neighboring areas of old cucurbit patches, under fallen leaves, in hedgerows, near their wild food sources (goldenrods and asters), or in garden debris. Adults emerge in early spring before cucurbits are available as food, and feed on pollen, petals, and leaves of alternative hosts.

Once cucurbits begin to emerge or once protective cover is removed, the adults migrate to their preferred hosts and begin chewing on leaves and stems. The beetles mate at this time and females deposit their eggs at the base of host plants, below the ground surface. Upon hatching (8-10 days) the larvae migrate to the root system and feed upon roots for 2-6 weeks, during which time they may consume the entire root system. Larvae pupate in the soil, emerging as adults in about one week. In fall, the adults return to the outlying areas for winter. Depending on geographic region and weather conditions, New Hampshire gardens may experience one to three generations of striped cucumber beetle in any given season.



Adult striped cucumber beetle. Credit: Alan T. Eaton.

The striped cucumber beetle is responsible for the spread of plant diseases such as bacterial wilt and cucumber mosaic.



Immature striped cucumber beetles in cantaloupe rind. Credit: Whitney Cranshaw, Colorado State University, Bugwood.org.



Damage by the striped cucumber beetle on melon (larval tunneling at crown and adult scarring). Credit: Whitney Cranshaw, Colorado State University, Bugwood.org.

Management

IPM Strategies:

- Cultural practices — Rotate cucurbit crops to a new place in the garden or farm each year. Cover planting beds with floating row covers immediately after planting seeds or setting out plants. Remove row covers as plants begin to bloom, to ensure adequate bee pollination. Applying a heavy mulch of straw, leaves, or grass clippings around established plants may help reduce striped cucumber beetle attacks.
- Sanitation — At the end of the season, destroy or bury crop debris (vines, leaves, remaining fruits) to deprive adults of overwintering spots. Consider destroying asters, goldenrod or other nearby pollen (food) sources for the adults.
- Monitoring — Early in the growing season, check plants frequently for adult beetles or chewing damage on cucurbit stems and foliage. Adult beetles are difficult to handpick, as they fly away or drop to the soil to hide when plants are disturbed. The presence of adult beetles or visible damage to plants may indicate a need for chemical control. The most susceptible time is from crop emergence or transplanting until the plant has reached the 5-leaved stage.
- Chemical Control — Hundreds of products are registered to control striped cucumber beetles. Check the labels of any product you purchase to ensure that both the target insect (striped cucumber beetle) and the plants you intend to spray are listed. Follow all label instructions. If the crop is in bloom when treatment is required, select an insecticide that is low risk to pollinators and spray at dusk. Avoid spraying insecticide in the morning on squash-family crops in bloom.

Guidelines for control of the striped cucumber beetle are in the [New England Vegetable Management Guide](#). For a more personalized recommendation, consult your county Agricultural Field Specialist.

Did You Know?

Do not use insecticides while cucurbit crops are in bloom. Most insecticides are hazardous to bees. An exception is surround, which do not kill insects.



Adult striped cucumber beetles feeding on cull pumpkin. Credit: Whitney Cranshaw, Colorado State University, Bugwood.org.

Summary

Table 1 summarizes key information on the striped cucumber beetle.

Table 1: Summary

Summary Table	
Damaging Stage	Adults, larvae
Part of Plant Attacked	Roots, leaves, and fruits
Overwintering Stage	Adults
Number of Generations per Year	One
Time of Year of Greatest Damage	May-September
Number of Pesticide Applications for Control	Zero to three

Notes: Refer to the text for more information on the striped cucumber beetle.

Stop! Read the label on every pesticide container each time before using the material. Pesticides must be applied only as directed on the label to be in compliance with the law. All pesticides listed in this publication are contingent upon continued registration. Contact the Division of Pesticide Control at (603) 271-3550 to check registration status. Dispose of empty containers safely, according to New Hampshire regulations.

Updated: July 2016 (Dr. Alan T. Eaton and Rachel Maccini)
Reformatted: July 2016 (Marie-Eve Jacques)

Visit our website:
extension.unh.edu

UNH Cooperative Extension brings information and education into the communities of the Granite State to help make New Hampshire's individuals, businesses, and communities more successful and its natural resources healthy and productive. For 100 years, our specialists have been tailoring contemporary, practical education to regional needs, helping create a well-informed citizenry while strengthening key economic sectors.

The University of New Hampshire Cooperative Extension is an equal opportunity educator and employer. University of New Hampshire, U.S. Department of Agriculture and New Hampshire counties cooperating.

About the Author

Dr. Alan T. Eaton is an Extension Specialist in Entomology and a professor at the University of New Hampshire. Much of his work is on management of fruit pests and ticks.

For More Information

State Office

Taylor Hall
59 College Rd.
Durham, NH 03824
<http://extension.unh.edu>

Education Center and Information Line

answers@unh.edu
1-877-EXT-GROW
(1-877-398-4769)
9 am–2 pm M–F
Search key words:
“UNH Education Center”