

FLINT POND  
OF HOLLIS  
LAKES LAY MONITORING PROGRAM  
1986

Freshwater Biology Group (FBG)  
University of New Hampshire  
Durham


by

Jeffrey A. Schloss

Coauthored and edited by

A.L. Baker

J.F. Haney



LAKES LAY MONITORING PROGRAM

To obtain more information about the Lakes Lay Monitoring Program (LLMP) contact the LLMP Coordinator (J. Schloss) at (603)-862-3848, Dr. Baker at 862-3845 or Dr. Haney at 862-2106.

## TABLE OF CONTENTS

ACKNOWLEDGEMENTS .....	iii
PROGRAM DESCRIPTION .....	v
The Lakes Lay Monitoring Program .....	v
COMMENTS AND RECOMMENDATIONS .....	vii
INTRODUCTION .....	1
Importance of long term monitoring .....	1
METHODS OF THE FRESHWATER BIOLOGY GROUP .....	5
Field and Laboratory Methods .....	5
Data analysis .....	10
RESULTS AND DISCUSSION OF FBG DATA .....	13
Water Transparency .....	13
Chlorophyll a .....	14
Dissolved Color .....	15
Total Phosphorus .....	15
Specific Conductivity .....	16
Stratification in the Deep Water Site .....	16
Dissolved Oxygen and Free CO <sub>2</sub> .....	16
pH .....	17
Alkalinity .....	20
Phytoplankton .....	20
Zooplankton .....	21
REFERENCES .....	23
APPENDIX	
Lay Monitor Data (1985) .....	A-1



## ACKNOWLEDGEMENTS

This was the second year of participation in the Lakes Lay Monitoring Program (LLMP) for the Flint Pond Association. The Lay Monitor Contact is Mr. Clement Deschenes and Alfred Harrison provided the boat for sampling. We encourage these and other interested members of the Flint Pond Association to continue monitoring during the 1987 season. We would also like to acknowledge Edwin Howes of the Nashua Regional Planning Commission for his dedication to the organization and maintenance of the LLMP for Flint Pond. Most important was the contribution by the Town of Hollis, which is a welcome demonstration of the support this monitoring program receives from the Town Selectmen.

The Freshwater Biology Group (FBG) is co-supervised by Dr. Alan Baker and Dr. James Haney. Members of the FBG summer field team included Tracy Kenealy, Jeff Schloss, Patricia McCarthy, Lori Sommer, Steve Thomas and Zhanyang Guo. Tracy and Jeff shared coordination of the program and were responsible for arranging the field trips, training lay monitors, and supervising the research team. Patricia and Lori were responsible for the preparation of chemical solutions, chlorophyll analysis and data entry. Steve was responsible for phosphorus chemistry and analysis. All team members participated in field work and chemical analyses. In

the fall, Alice Hibberd assisted in data organization and data entry and Jeff continued as **LLMP** Coordinator responsible for data interpretation and report writing.

The **FBG** would like to thank the University of New Hampshire Undesignated Gifts Committee for the partial funding of the coordinator position. Eileen Wong of the Department Zoology provided accounting and secretarial service. The Department of Botany and Plant Pathology provided lab and storage space. We would also like to recognize the UNH Office of Computer Services for the provision of computer time and data storage space.

Participating groups in the **LLMP** for 1986 included: The New Hampshire Audubon Society, Derry Conservation Commission, Nashua Regional Planning Commission, Center Harbor Bay Conservation Commission, Governor's Island Club Inc., Little Island Pond Rod and Gun Club, Walker's Pond Conservation Society, United Associations of Alton, the associations of Baboosic Lake, Beaver Lake, Berry Bay, Bow Lake Camp Owners, Lake Chocorua, Flint Pond, Lake Kanasatka Watershed, Langdon Cove, Long Island Landowners, Moultonbouro Bay, Lake Winnipесаaukee, Naticook Lake, Newfound Lake, Nippo Lake, Scruton Pond, Silver Lake (Hollis), Silver Lake (Madison), Squam Lake, Sunset Lake, Lake Winona, and Lake Wentworth and the towns of Alton, Amherst, Hollis, Merrimack and Stratham.

## PROGRAM DESCRIPTION

### The Lakes Lay Monitoring Program

The New Hampshire Lakes Lay Monitoring Program (LLMP) is a research and educational function of the Freshwater Biology Group (FBG) at the University of New Hampshire co-directed by Professors Alan Baker, Department of Botany and Plant Pathology, and James Haney, Department of Zoology and coordinated by Jeffrey Schloss. The program involves the cooperative participation of lake residents, lake associations, conservation and planning commissions and local governments with University faculty and students. Developed in 1978 around Squam Lake, the program has grown to include more than 30 lakes throughout New Hampshire.

As a research project, the LLMP has investigated the extent of lake degradation caused by perturbations such as acid rain, septic and agricultural runoff, and lakeshore development. Essentially the monitors in the program collect data once each week. The data are stored on a computer, the results are analyzed periodically, and interpretive reports are written that include graphics and statistical analyses. A major goal is to detect any short or long-term changes in the water quality of the lakes. To that end a long-term data base has been established.

As an educational tool, several students are trained each year to collect and analyze lakewater samples for physical, chemical and biological parameters, and to interpret water quality data. In addition, more than 200 "lay" monitors have been trained to monitor their own lakes and educated about lake water quality.

As a service to the state and to local communities, the reports of the LLMP are available at cost, and should prove useful to lake residents, conservationists, developers and land-use planners. Also, LLMP staff members conduct workshops, lectures and informal talks on various lake related topics and hold advisory positions on many municipal and private conservation and planning boards. The LLMP is a not-for-profit organization with funding derived primarily from the participating groups.

## COMMENTS AND RECOMMENDATIONS

1) We recommend that each association, including the Flint Pond Association continue to develop their data base on lake water quality through continuation of the long term monitoring program. The data base will provide information on the short and long-term cyclic variability that occurs in the lake and eventually will enable more reliable predictions of water quality trends. The database was initiated in 1985 but no monitoring by lay persons was undertaken in 1986.

2) We recommend phosphorus testing to be taken during the summer months. As early as possible for the initial sampling combined with sampling of the lake during a times of heavy use (ie: 4 July, Labor Day) or late in the season when septic systems have been put through a full seasons use.

3) The **FBG** trip provided a more in-depth analysis of the lake during late summer conditions. We recommend one or more **FBG** trips in 1987.

4) We invite the Flint monitors to participate in our preliminary investigation of the effect of boat traffic on lakes. All that would be required is sampling in the morning and late in the afternoon on a "quiet day" followed by the same sampling approach on a day of heavy boat traffic. A discount for sample processing will be offered to try to



minimize costs of additional testing. Contact the coordinator for further information.

5) As a general addition to our Lakes Lay Monitoring Program, we recommend that each lake in the Program begin monitoring the condition of the fish taken from the lake. The "Fish Monitoring" will require at least one lay monitor to record the species, length and weight and collect a sample of fish scales for each fish examined. In most lakes this will involve periodic creel census of sport fishermen on the lake. The required equipment, supplies and analytical costs will be approximately \$100. Explanation of procedures and fish identification will be given to monitors who decide to measure this parameter.

Length-to-weight ratios give a measure of the nutritional condition of the fish. Age analysis of the fish scales (to be done at UNH) will tell how old each fish is. Together, these variables can help to track changes in the condition of the fish populations in the lake, and, of course, the "health" of the lake.

## INTRODUCTION

### Importance of long term monitoring

A major goal of this program is to identify any short or long-term changes in the water quality of the lakes. Of major concern, is the detection of cultural eutrophication; increases in the productivity of the lake due to the addition of nutrients from human activities. Changes in the natural buffering capacity of the lakes in the program is also a topic of great concern since New Hampshire receives large amounts of acid precipitation. Weekly sampling of a lake during a single summer provides information only on the variation that occurs. Short-term differences may be due to variations in weather or lake activity, or other chance events. The resulting short-term fluctuations may be unrelated to the actual long-term trend.

As an example, a 30 year study of a lake may indicate a long-term trend toward eutrophy (Fig. 1). Yet if only the data from a five year period (ie: Fig 1, years 1975-80) are examined, no apparent trends can be seen. If only two years are examined, the data suggest a decrease in eutrophy! Monitoring carried out weekly over the course of many summers can provide the information required to distinguish between short-term fluctuation ("noise") and long-term

trends ("signal"). To that end, each lake must establish a long-term data base.




Figure 1. Eutrophication of a hypothetical lake over time. Circled area is enlarged for comparison between short and long-term trends.

The number of seasons it takes to discern between the noise and the signal is not the same for each lake. Evaluation and interpretation of a long term data base will indicate that the water quality of the lake has worsened, improved, or remained the same. As more data is collected prediction of current and future trends can be made. No matter what the outcome, this information is essential for the intelligent management of the lake.

There are also short-term uses for lay monitoring data. The examination of different stations in a lake can disclose specific problems and corrective action can be initiated to handle the situation before it becomes more serious. On a lighter note, some associations post their weekly data for use in determining the best depths for finding fish!

It takes a considerable amount of effort as well as a deep concern for one's lake to be a lay monitor. Many times a monitor has to brave inclement weather or heavy boat traffic to collect samples. Sometimes it even may seem that one week's data is just the same as the next. Yet every sampling provides important information on the variability of the lake.

Every data sheet the **LLMP** receives is significant to further the understanding of the lakes in the program. We are pleased with the interest and commitment of our lay monitors and are proud that their work is what makes the **LLMP** the most extensive, and we believe, the best volunteer program of its kind.



## METHODS OF THE FRESHWATER BIOLOGY GROUP

The Freshwater Biology Group (FBG) research team took one trip to Flint Pond and conducted several tests which included measurements of sunlight penetration into the water, dissolved oxygen, temperature, alkalinity, free (unbound) carbon dioxide, pH, specific conductivity, chlorophyll a, dissolved color, total phosphorus, and a survey of the microscopic plants (phytoplankton) and animals (zooplankton). The input, storage and analysis of all LLMP data is also the responsibility of the FBG.

### Field and Laboratory Methods

On the lake, a dissolved oxygen and temperature profile was taken using a Yellow Springs Instruments Model 54A Oxygen/Temperature meter with a submersible probe. Readings were taken at one-meter intervals throughout the epilimnion and hypolimnion, and at one-half meter intervals through the metalimnion.

Sunlight and skylight penetration into the water was measured with a Whitney submersible photometer model LMA-8A, off the sunny side of the boat. The coefficient of light extinction was calculated from the relative light intensities measured.

Water clarity was measured by lowering a secchi disk (approximately 20 cm. or 8 inches) through the water off the

shaded side of the boat, and noting the average of the depths at which it disappeared upon lowering and reappeared when being raised (the cord attached to the secchi disk is marked in one tenth of a meter for the first half meter and in one-half meters thereafter). Water clarity was determined while holding a view-scope just below the surface to eliminate effects of surface reflection and wave action. This was repeated two or three times, and an average to the nearest one-tenth of a meter was recorded.

Samples of lake water chemistry to be analyzed for dissolved oxygen, alkalinity, free (unbound) carbon dioxide, pH, and specific conductivity were collected with a 3-liter Van Dorn bottle at depths which represented the surface, mid-epilimnion, metalimnion, and hypolimnion. Alkalinity, free carbon dioxide, and pH samples were stored on ice in 250 milliliter polyethylene bottles and were analyzed in the field within 1 to 2 hours of sampling. Specific conductivity samples were analyzed in the FBG lab at room temperature.

In addition to the oxygen profile taken, the dissolved oxygen (DO) concentration of specific lakewater samples (epilimnetic and hypolimnetic) were determined chemically with the azide modification of the Winkler method (EPA 1979). The precision of the method provides a standard for the electronic probe. Water is collected in 350 ml biological oxygen demand (BOD) bottles and fixed with two

reagents, manganese sulfate and alkali-iodine-azide. A loose precipitate (floc) of manganic hydroxide is formed that is equivalent to all dissolved oxygen originally present in the sample. Concentrated sulphuric acid is added to the bottle which causes a stoichiometric release of dissolved iodine equal to the original amount of dissolved oxygen present. A known quantity of sample is then titrated to an equivalence point using .0250N phenylarsine oxide titrant (similar to, but more stable than, sodium thiosulphate which may also be used) and a starch indicator solution. The end-point is reached when the purple colored iodine-starch complex is reduced and the solution becomes colorless. The amount of titrant added is recorded to the nearest 0.1 ml and concentrations are reported to the nearest 0.2 milligrams dissolved oxygen per liter.

To determine the alkalinity, lake water samples were titrated with 0.002 N sulphuric acid in the presence of the indicator methyl red/bromocresol green to a pH of 5.1 (grey endpoint) and 4.6 (pink endpoint). The amount of titrant used (dilute sulphuric acid) was recorded to the nearest 0.1 ml, equivalent to milligrams of calcium carbonate per liter. Values reported can be converted to microequivalents of calcium carbonate using a multiplication factor of 20.

"Free" carbon dioxide concentration was determined by titrating the fresh lakewater samples with 0.0027 N sodium



hydroxide to a final endpoint pH of 8.3, in the presence of the indicator dye phenolphthalein.

Lakewater pH was measured with a digital pH meter (Beckman model phi 44 ) equipped with a combination probe (Orion Co.) and an automatic temperature compensating probe. The meter was calibrated with pH 4 and pH 7 buffer solutions and then the probe was allowed to equilibrate in the lake water for at least thirty minutes prior to sample analysis.

Specific conductivity was measured with a Barnstead Conductivity Bridge Model PM-70CB , with a model B-10 probe (cell constant = 1.0). Corrections were made for sample temperatures with a standard curve of potassium chloride solution conductivity versus temperature. Results are reported as micro-Siemens (uS; where uS equals  $\mu\text{mho cm}^{-2}$ ) standardized to 18<sup>o</sup> C.

Samples to be analyzed for chlorophyll a, total phosphorus, and phytoplankton were collected with a vertical tube sampler into a 2.5 liter dark plastic bottle. Chlorophyll samples were filtered through a 0.45 micron membrane filter and air-dried in the dark until analysis. The chlorophyll a content was analyzed by extracting the chlorophyll with a 95% acetone solution saturated with magnesium carbonate. The samples were then centrifuged and their light absorbance read at two standard wavelengths (663 and 750 nanometers) with a Baush and Lomb model 710

spectrophotometer equipped with 50mm cuvettes. An absorptivity value of  $84 \text{ gm liter}^{-1} \text{ cm}^{-1}$  (Vollenweider 1969) was used for calculating the concentrations.

Dissolved color samples of the filtrate from **FBG** and lay monitor chlorophyll filtrations was determined by reading the absorbance of the samples at two different wavelengths (440 and 493 nanometers) in a 50mm light path. The two readings were converted to the more widely used platinum cobalt color values (ptu) using standard curves of the absorbance of chloroplatinate.

Phosphorus samples were preserved with 1.0 milliliter of concentrated sulphuric acid and refrigerated until analysis. Also, phosphorus samples from lay monitors were received by the **FBG** in a refrigerated or frozen state, and stored cold until analysis. To determine the total phosphorus content, ammonium persulfate and 11 N sulfuric acid was added to digest the total phosphorus, and the samples were autoclaved for thirty minutes at 250 to 260 degrees C. Reagents included potassium antimony tartrate, ammonium molybdate, and a solution of ascorbic acid mixed fresh before each sample run (E.P.A. 1979). Absorbance of the blue phosphorus complex was measured with a spectrophotometer at 650 nanometers. A standard curve of the absorbance of a potassium phosphate (monobasic) solution to convert the readings to total phosphorus concentrations.

Each sample was analyzed twice and an average of the two values taken as the phosphorus content in parts per billion (ppb).

Phytoplankton samples were preserved with iodine (Lugol's solution) immediately after collection. Algae were later identified and counted with an inverted microscope after settling for 24 hours in 5 or 10 ml counting chambers. At least 200 individual algal "units" were counted with a modified scan technique (Baker, 1973). Phytoplankton are reported to species level whenever possible.

Zooplankton samples were collected with a plankton net (30 centimeter diameter, 150 micron porosity) towed vertically through the oxygenated portion of the water (>0.5 ppm oxygen). Samples were immediately preserved in a 4% formalin-sucrose solution (Haney and Hall, 1973). Organisms were identified to species whenever possible. Subsampling, whenever necessary, was done with a 1 ml Hensen-Stemple pipette. Repeated subsamples were analyzed until at least 100 organisms were counted.

#### Data analysis

Incoming data are received through the mail during the sampling season and are first filed in an "incoming data" book. This provides temporary storage until the corresponding chlorophyll and/or phosphorus sample for each data sheet is analyzed. All data, including date, lake,

site, secchi disk depth, chlorophyll a and phosphorus concentrations, alkalinity, and color measurements, are filed and stored on the FBG computerized data-management system that utilizes a mainframe DEC VAX-8650 computer and an IBM compatible microcomputer (Zenith Data Systems 158). With full use of relational data bases, such as S1032 and Dbase III+ data can be easily retrieved by lake, date, station or by parameter and used for individual reports and program summaries for each year.

Statistical treatment of the data from each lake, produced for level III reports, includes a comparison of seasonal tendencies found throughout the year, monthly means for the different parameters tested, and confidence levels for each site. The same comparisons are made on a yearly basis if the lake has been in the program for two years or more. Where sufficient data are available from several years, regression analyses and other statistical tests can be performed. Such analyses may identify trends and help explain variations in the data (eg. secchi disk depth, chlorophyll a, color). In addition, data from a lake may be compared with other lakes in the program, other computerized data bases (New Hampshire Water Supply and Pollution Control, New Hampshire Fish and Game, EPA Surface Water Survey and others) and to published water quality classifications.

Trophic boundaries of Forsberg and Ryding (1980) of transparency, chlorophyll a, and total phosphorus are used as criteria in discussions of the trophic state of the program lakes. Phytoplankton are reported both as species and classes. Crustacean zooplankton were classified into one of four categories depending on their size (large or small) and their feeding preferences (herbivore or predator) with a modified version of criteria from Sprules (1980). The differences in abundance between the different groups allow for a more complete description of the zooplankton community and the trophic classification of lakes.

## RESULTS AND DISCUSSION OF FBG DATA

Sampling of Flint Pond by the FBG field team took place on 22 August at site 1 Deep. This site is located at the north end of the lake and is approximately 2.5 meters deep. Table 1 presents the findings from all FBG trips to date. Lay monitor data is summarized in the appendix.

TABLE 1 RESULTS OF FBG INTEGRATED SAMPLING FOR FLINT POND

DATE	Secchi Depth (meters)	Chlorophyll (mg m <sup>-3</sup> )	Color (ptu)	Total Phosphorus (ppb)
7/02/85	B.O.*	1.0	40	16
7/11/85	2.0	2.9	26	12
8/02/85	2.3	7.4	63	10
8/23/85	B.O.	1.5	23	4
8/22/86	2.0	1.7	41	11

\*B.O. = secchi disk visible on bottom.

### Water Transparency

Secchi Disk depth is a measure of the water transparency. The deeper the depth of secchi disk disappearance, the more transparent the lake water; light penetrates deeper if there is little dissolved and/or particulate matter (which includes both living and non-living particles) to absorb and scatter it.

One must be cautious when interpreting transparency measurements for a shallow lake. Mixing in the water column due to wind or boat traffic will bring up sediment from the bottom which will decrease transparency. A transparency of less than 4.0 meters is typical of more productive lakes while values below 2.5 are common in very productive lakes. Previous lay monitoring data in 1985 at a deeper site had a secchi depth range of 3.3 to 4.5 meters.

### Chlorophyll a

The chlorophyll a concentration is a measurement of the standing crop of phytoplankton and is often used to classify lakes into categories of productivity called trophic states. **Eutrophic** lakes are highly productive with large amounts of algae and aquatic plants due to nutrient enrichment. **Oligotrophic** lakes have low productivity and low nutrient levels and **mesotrophic** lakes are intermediate in productivity.

Chlorophyll concentration in 1986 at the end of August was moderate as in the previous years sampling by the FBG and the lay monitors. The high concentration during early August 1985 could have been a bloom event or a result of the stirring up of the bottom waters.

### Dissolved Color

The dissolved color of lakes is generally due to dissolved organic matter from humic substances, which are naturally-occurring polyphenolic compounds leached from decayed vegetation. Highly colored or "stained" lakes have a "tea" color. Such substances generally do not threaten water quality except as they diminish sunlight penetration into deep waters.

Dissolved color was high at site 1. To put the color level in perspective, dissolved color concentrations of all lakes participating in the LLMP in 1986 were in the range <1 to 117 ptu with an average of 18.5. Thus, Flint Pond color was above average compared to all lakes in the program.

### Total Phosphorus

Of the two "nutrients" most important to the growth of aquatic plants, nitrogen and phosphorus, it is generally observed that phosphorus is the more limiting to plant growth, and therefore the more important to monitor and control. Phosphorus is generally present in lower concentrations, and its sources primarily originate from anthropogenic activity in a watershed. Nitrogen can be fixed from the atmosphere by many bloom-forming blue-green bacteria, and thus it is difficult to control. The total phosphorus includes all dissolved phosphorus as well as



phosphorus contained in or adhered to suspended particulates such as sediment and plankton.

Phosphorus concentration on 22 August was 11 ppb at site 1. Concentrations of total phosphorus at this site during past FBG trips has generally approached mesotrophic levels (greater than 15 ppb).

#### Specific Conductivity

The specific conductance of a water sample indicates concentrations of dissolved salts. Leaking septic systems and de-icing salt runoff from highways can cause high conductivity values. Conductivity values at Flint Pond in 1986 had a range of 158 to 163  $\mu\text{S}$  suggesting that the lake is receiving road salt runoff and/or septic input near site 1.

#### Stratification in the Deep Water Site

Profiles of temperature for site 1 did not indicate the typical pattern of temperature stratification where a layer of warmer water (the epilimnion) overlies a deeper layer of cold water (hypolimnion). The layer that separates the two regions characterized by a sharp drop in temperature with depth is called the thermocline or metalimnion.

#### Dissolved Oxygen and Free CO<sub>2</sub>

Oxygen concentration was high in the epilimnion and decreased to very low concentrations in the bottom waters


(Fig. 1). Carbon dioxide in bottom waters is moderate. Note the increase just above the bottom sediment. Carbon dioxide is generated and can accumulate in aquatic systems as a result of the respiration of a wide variety of organisms in the water. Plants (including the phytoplankton) take up free carbon dioxide and produce oxygen during the day, but respire at night along with the aquatic animals and bacteria. Carbon dioxide usually accumulates in the bottom waters of more productive systems where large amounts of organic material, produced within and around the lake, support large bacterial populations. Breakdown of organic matter, respiration and fermentation by the bacteria in the water and sediments consumes oxygen and releases carbon dioxide. Increases in dissolved carbon dioxide result in the decrease of the lakewater pH.

#### pH

The pH is a way of expressing the acidic level of lake water, and is measured with an electrical probe sensitive to hydrogen ion activity. The pH scale has a range of 1 (very acidic) to 14 (very "basic" or alkaline) and is logarithmic (ie: changes in 1 pH unit reflect an order of magnitude, ie: 10 times, difference in hydrogen ion concentration). Most

Figure 1 - Profiles of dissolved oxygen (O<sub>2</sub>), "free" carbon dioxide (CO<sub>2</sub>) and pH on 22 August 1986 at Flint Pond site 1 Deep. Parameters are scaled 0 to 8 by the uppermost horizontal axis. Oxygen was measured at one-half meter intervals, other parameters were sampled at the discrete depths indicated by the x's and asterisks. Water temperature was constant.

FLINT POND 1 DEEP 22 August 1986



DISSOLVED GASES AND pH

aquatic organisms tolerate a limited range of pH and most fish species require a pH of 5.5 or higher for successful growth and reproduction.

Surface pH was 6.9 at site 1. The range of surface water pH for all LLMP lakes was 5.2 to 7.2. The pH increased with depth to 7.2 (Fig. 1). At present the pH of Flint Pond is within the optimum range of many aquatic organisms.

### Alkalinity

Alkalinity is a measure of the buffering capacity of the lake water. The higher the value the more acid that can be neutralized. Typically lakes in New Hampshire have low alkalinities due to the absence of carbonates and other natural buffering minerals in the bedrock of lake watersheds.

Surface alkalinity at site 2 was 33.9 mg CaCO<sub>3</sub> liter<sup>-1</sup> and increased to 34.1 mg CaCO<sub>3</sub> liter<sup>-1</sup> at 2.0 meters. The alkalinity at Flint Pond is high compared to most lakes in New Hampshire and is the second only to Beaver Lake in Derry for all lakes in the LLMP in 1986.

### Phytoplankton

Concentration of phytoplankton at site 1, was moderate to high, 4176 organisms per ml, typical of open-water plankton in lakes of intermediate productivity. Unlike the

previous years community, the golden algae (Chrysophyceae) specifically an unidentified flagellate was dominant and the blue-green bacteria (Cyanophyceae) were subdominant.

### Zooplankton

The macrozooplankton (caught using a 150 micrometer mesh net; excluding copepod nauplii) in the oxygenated waters of site 1 numbered 72.4 organisms per liter. This is a high value. Calanoid copepods were the dominant herbivores ("phytoplankton eaters") with a concentration of 34.0 organisms per liter. The herbivorous cladoceran Bosmina was subdominant. As in 1985 small herbivores predominated. Bosmina and the predatory cyclopoid copepods were at the highest concentration ever reported so far for Flint Pond.



## REFERENCES

- American Public Health Association. (APHA) 1975. Standard Methods for the Examination of Water and Wastewater 14th edition. APHA, AWWA, WPCF. Washington, D.C.
- Baker, A.L. 1973. Microstratification of phytoplankton in selected Minnesota lakes. Ph. D. thesis, University of Minnesota.
- Carlson, R.E. 1977. A trophic state index for lakes. *Limnol. Oceanogr.* 22:361-379.
- Edmondson, W.T. 1937. Food conditions in some New Hampshire lakes. In: Biological survey of the Androscoggin, Saco and coastal watersheds. (Report of E.E. Hoover.) New Hampshire Fish and Game Commission, Concord, New Hampshire.
- Forsberg, C. and S.O. Ryding. 1980. Eutrophication parameters and trophic state indices in 30 Swedish wastewater receiving lakes. *Arch. Hydrobiol.* 89:189-207
- Gallup, D.N. 1969. Zooplankton distributions and zooplankton-phytoplankton relationships in a mesotrophic lake. Ph.D. Thesis, University of New Hampshire.
- Haney, J.F. and D.J. Hall. 1973. Sugar-coated Daphnia: a preservation technique for Cladocera. *Limnol. Oceanogr.* 18:331-333.
- Hoover, E.E. 1936. Preliminary biological survey of some New Hampshire lakes. Survey report no. 1. New Hampshire Fish and Game Department, Concord, New Hampshire.
- Hoover, E.E. 1937. Biological survey of the Androscoggin, Saco, and coastal watersheds. Survey report no. 2. New Hampshire Fish and Game Department, Concord, New Hampshire.
- Hoover, E.E. 1938. Biological Survey of the Merrimack watershed. Survey report no. 3. New Hampshire Fish and Game Department, Concord, New Hampshire.
- Hutchinson, G.E. 1967. A treatise on limnology, vol. 2. John Wiley and Sons, New York.
- Lind, O.T. 1979. Handbook of common methods in limnology. C.V. Mosby, St. Louis.
- Lorenzen, M.W. 1980. Use of chlorophyll-Secchi disk relationships. *Limnol. Oceanogr.* 25:371-372.



- New Hampshire Water Supply and Pollution Control Commission. 1981. Classification and priority listing of New Hampshire lakes. Staff report no. 121. Concord, New Hampshire.
- Newell, A.E. 1970. Biological survey of the lakes and ponds in Cheshire, Hillsborough and Rockingham Counties. Survey report no. 8c. New Hampshire Fish and Game Department, Concord, New Hampshire.
- Newell, A.E. 1972. Biological survey of the lakes and ponds in Coos, Grafton and Carroll Counties. Survey report no. 8a. New Hampshire Fish and Game Department, Concord, New Hampshire.
- Newell, A.E. 1977. Biological survey of the lakes and ponds in Sullivan, Merrimack, Belknap and Strafford Counties. Survey report no. 8b. New Hampshire Fish and Game Department, Concord, New Hampshire.
- Schindler, D.W., et al. 1985. Long-term ecosystem stress: Effects of years of experimental acidification on a small lake. *Science*. 228:1395-1400.
- Sprules, W.G. 1980. Zoogeographic patterns in size structure of zooplankton communities with possible applications to lake ecosystem modeling and management. in W.C. Kerfoot ed. *Evolution and Ecology of Zooplankton Communities*. University Press of New England. Dartmouth. pp642-656.
- U.S. Environmental Protection Agency. 1979. A manual of methods for chemical analysis of water and wastes. Office of Technology Transfer, Cincinnati. PA-600/4-79-020.
- Vollenweider, R.A. 1969. A manual on methods for measuring primary productivity in aquatic environments. International Biological Programme. Blackwell Scientific Publications, Oxford.
- Wetzel, R.G. 1983. *Limnology*. Saunders College Publishing, Philadelphia.

Flint Lake Data on file as of 05/03/1987

Lakes Lay Monitoring Program, U.N.H.

[Lay Monitor Data]

Flint Lake

-- subset of trophic indicators, all sites, 1985

1985 SUMMARY

Average transparency:	3.9	(1985:	16	values)
Average chlorophyll:	1.4	(1985:	12	values)
Average phosphorus:	21.4	(1985:	17	values)
Average alk (gray):	24.9	(1985:	16	values)
Average alk (pink):	26.2	(1985:	16	values)

SITE	DA	SDD	CHL	TP	ALKG	ALKP	COLOR
Devlin Bch	08/14/1985	---	---	4.6	---	---	---
Devlin Bch	08/29/1985	---	---	9.4	---	---	---
Flint A	06/17/1985	---	---	16.5	---	---	---
Flint A	08/06/1985	3.4	0.7	---	27.5	27.6	---
Flint A	08/14/1985	3.3	0.4	3.5	28.1	29.9	---
Flint A	08/21/1985	4.0	1.9	---	27.1	27.8	30.7
Flint A	08/29/1985	3.7	0.6	---	22.1	22.3	---
Flint A	09/04/1985	4.2	0.4	---	25.0	27.4	---
Flint A	09/14/1985	4.5	1.6	---	28.2	29.9	---
Flint A	09/18/1985	4.5	---	---	2.7	2.8	30.7
Flint A	10/01/1985	4.5	---	---	29.1	29.9	---
Flint B	06/17/1985	---	---	9.1	---	---	---
Flint B	08/06/1985	3.8	1.9	---	26.4	27.3	---
Flint B	08/14/1985	3.6	2.4	4.9	28.3	30.0	---
Flint B	08/21/1985	3.5	2.6	---	26.4	27.1	25.5
Flint B	08/29/1985	3.4	0.7	---	25.0	28.2	---
Flint B	09/04/1985	4.0	1.9	---	27.4	29.9	---
Flint B	09/14/1985	4.0	1.6	---	25.8	27.4	---
Flint B	09/18/1985	4.1	---	---	25.8	27.4	33.2
Flint B	10/01/1985	4.0	---	---	24.2	25.0	---
Flint D1	06/17/1985	---	---	73.5	---	---	---
Flint D2	06/17/1985	---	---	17.5	---	---	---
Flint D3	06/17/1985	---	---	103.0	---	---	---
Flint E	06/17/1985	---	---	68.5	---	---	---
Flint E	08/14/1985	---	---	5.9	---	---	---
No Label!	08/14/1985	---	---	6.4	---	---	---
No Label!	08/14/1985	---	---	9.4	---	---	---
No Label!	08/29/1985	---	---	12.1	---	---	---
No Label!	08/29/1985	---	---	6.6	---	---	---
No Label!	08/29/1985	---	---	7.1	---	---	---
No Label!	08/29/1985	---	---	5.4	---	---	---