Lake Wentworth

WENTWORTH LAKE

1992

LAKES LAY MONITORING PROGRAM

by

Jeffrey A. Schloss

Robert K. Craycraft

edited by

A.L. Baker and J.F. Haney

NEW HAMPSHIRE LAKES LAY MONITORING PROGRAM



NH LLMP

FRESHWATER BIOLOGY GROUP

University of New Hampshire

Durham

UNIVERSITY OF NEW HAMPSHIRE

COOPERATIVE EXTENSION

Helping You To Put Knowledge And Research To Work

To obtain more information about the NH Lakes Lay Monitoring Program (NH LLMP) contact the Coordinator (J.Schloss) at (603) 862-3848 Dr. Baker at 862-3845 or Dr. Haney at 862-2106

PARAMETERS SAMPLED
NH LAKES LAY MONITORING PROGRAM



PREFACE

This report contains the findings of a water quality survey of Wentworth and Crescent Lakes, Wolfeboro, New Hampshire, conducted in the summer of 1992 by the **Freshwater Biology Group (FBG)** of the University of New Hampshire and the Lake Wentworth Association.

The report is written with the concerned lake resident in mind and contains a brief, non-technical summary of 1992 results as well as more detailed "Introduction" and "Discussion" sections. Graphic display of data is included, in addition to listings of data in appendices, to aid visual perspective.

TABLE OF CONTENTS

PREFACE 1

TABLE OF CONTENTS 3

ACKNOWLEDGEMENTS 5

CRESCENT LAKE - 1992 NON-TECHNICAL SUMMARY 7

WENTWORTH LAKE - 1992 NON-TECHNICAL SUMMARY 11

COMMENTS AND RECOMMENDATIONS 14
- The New Hampshire Lakes Lay Monitoring Program 15
- Importance of Long-term Monitoring 15
- Purpose and Scope of This Study 17

DISCUSSION OF LAKE MONITORING MEASUREMENTS 19
- Thermal Stratification in the Deep Water Sites 19
- Water Transparency 19
- Chlorophyll a 20
- Dissolved Color 21
- Total Phosphorus 22
- pH * 23
- Alkalinity 23
- Specific Conductivity * 25
- Dissolved Oxygen and Free Carbon Dioxide * 25
- Underwater Light * 26
- Indicator Bacteria * 27
- Phytoplankton * 28
- Zooplankton * 29
- Fish Condition 30

REFERENCES 31

REPORT FIGURES 33

DATA A-1

LAKE DIAGRAMS B-1

GLOSSARY C-1

ACKNOWLEDGEMENTS

This was the ninth year of participation in the **Lakes Lay Monitoring Program (LLMP)** for the Wentworth Lake Association monitors. The Lay Monitor of Crescent lake was Gus G. Freygang while Lawrence Goldthwait, John Nichols and Alfred Flynn were the Lay Monitors of Wentworth Lake. The coordinator of the volunteer monitoring program and liaison to the **FBG** was John Nichols. The **Freshwater Biology Group (FBG)** congratulates the Lay Monitors on the quality of their work, and the time and effort put forth. The **FBG** would also like to acknowledge the late Dr. Richard Goldthwait, through whose efforts the volunteer monitoring program of Wentworth and Crescent Lakes became established and has continued to grow. We encourage interested members of the Wentworth Lake Association to continue monitoring during the 1993 season.

The **Freshwater Biology Group** is a not-for-profit research program co-supervised by Dr. Alan Baker and Dr. James Haney and coordinated by Jeffrey Schloss. Members of the **FBG** summer field team included Jeffrey Schloss, Robert Craycraft, Gregg Vereb, Gregg Stevens, Sean Proll, Matt Denneen and Robert Banks. Other **FBG** staff assisting in the fall were Eric Betke, Amanda Fifield, Jessica Chappell and Phil Lucason.

The **FBG** acknowledges the University of New Hampshire Cooperative Extension for funding and furnishing office, laboratory and storage space. The College of Life Sciences and Agriculture provided accounting support and the UNH Office of Computer Services provided computer time and data storage allocations.

Participating groups in the **LLMP** include: The New Hampshire Audubon Society, Derry Conservation Commission, Dublin Garden Club, Nashua Regional Planning Commission, Center Harbor Bay Conservation Commission, Governor's Island Club Inc., Little Island Pond Rod and Gun Club, Walker's Pond Conservation Society, United Associations of Alton, the Pemaquid Watershed Study Group, the associations of Baboosic Lake, Beaver Lake, Berry Bay, Big Island Pond, Bow Lake Camp Owners, Chesham

Pond, Lake Chocorua, Crystal Lake, Cunningham Pond, Dublin Lake, Glines Island, Goose Pond, Great East Lake, Lake Kanasatka Watershed, Langdon Cove, Long Island Landowners, Lovell Lake, Marchs Pond, Mascoma Lake, Mendum's Pond, Meredith Bay Rotary Club, Merrymeeting Lake, Milton Ponds Lake Lay Monitoring, Mirror Lake (Tuftonboro), Moultonbouro Bay, Lake Winnipesaukee, Naticook Lake, Newfound Lake, Nippo Lake, Peaporridge Pond, Perkins Pond, Pleasant Lake, Silver Lake (Hollis), Silver Lake (Harrisville), Silver Lake (Madison), Silver Lake (Tilton), Squam Lakes, Lake Sunapee, Sunset Lake, Lake Waukewan, Lake Winona, Wentworth Lake and the towns of Alton, Amherst, Enfield, Hollis, Madison, Merrimack, Strafford and Wolfeboro.

CRESCENT LAKE

1992 NON-TECHNICAL SUMMARY

Monitoring was undertaken at Crescent Lake by the volunteer monitor from June 16 to September 16. A more in-depth analysis was conducted by the FBG on August 3.

1) Water transparency at Crescent Lake was moderate to high, the sign of a relatively clear and unproductive lake. The secchi disk was visible as far down as 5.6 meters (18.2 feet) and the transparency average was 4.9 meters at Site 6 Center. This indicates the deepwater site on the lake is relatively low in dissolved color and suspended matter such as algae and particulates. Average water clarity increased in 1992 but remains below the average levels recorded in 1984 through 1990.

2) Chlorophyll *a* concentrations for the surface waters of Crescent Lake were low to moderate. Concentrations in the mixed layer of water averaged 3.0 milligrams per cubic meter (3.0 mg m^{-3}, equivalent to about 3.0 parts chlorophyll per billion parts water) at Site 6 Center. Generally, concentrations below 3 mg m^{-3} are common to less productive clear lakes and values above 7 mg m^{-3} are common in productive lakes. The 1992 average chlorophyll *a* level is indicative of a borderline unproductive (oligotrophic) to moderately productive (mesotrophic) lake and is similar to the average level observed in 1991. Average chlorophyll *a* levels in both 1991 and 1992 were below the historical average. However, this is most likely the result of the below average precipitation over that period, resulting in minimal nutrient loading to stimulate algal growth. With 1993 off to a more typical year in respect to precipitation, the lake is likely to respond by exhibiting higher chlorophyll *a* concentrations during the 1993 sampling season.

3) Dissolved lakewater color levels for Crescent Lake were low in 1992, 18.3 ptu (platinate color units), and less than the average levels of 26 ptu in other program lakes.

Small increases in water color from the natural breakdown of plant materials in and around a lake are not considered to be detrimental to water quality. However, increased color can lower water transparency, and hence, change the public perception of water quality. Large amounts of dissolved color may occur naturally but also occur during deforestation and development within the watershed. High color levels can actually mask the ability of the secchi disk transparency to predict chlorophyll levels.

4) Total phosphorus (nutrient) levels were low when sampled by the **FBG**, 2.0 parts per billion (ppb) on August 3 and low to moderate, 10.4 ppb, when sampled by the volunteer monitor on September 12. A concentration of 15 ppb is commonly thought of as the boundary between less productive and more productive lakes.

5) The pH of the surface waters of the lake, measured by the **FBG** and volunteer monitor, remains within the optimum range for most aquatic organisms. The alkalinity of the lake, the lake's ability to buffer acid input, is low, and slightly less than the average of 6.3 units for **LLMP** program lakes. While the alkalinity level is low, it remains sufficient to buffer against fluctuations in pH caused by acid loadings.

6) The specific conductivity of the deep site on Crescent lake was low, between 43.1 and 44.6 micro-Siemans. High conductivity values can indicate the presence of septic leachate or deicing salt runoff.

7) Temperature profiles collected by the volunteer monitor disclosed the typical temperature stratification patterns for shallow New Hampshire lakes. With the lake becoming only partially stratified when the weather was calm. The oxygen content of the bottom waters remained high, above 3 milligrams per liter, (the minimum concentration required for the successful growth and reproduction of most warm-water fish) to the lake-bottom when measured by the **FBG** in early August.

8) For all measurements considered and averaged for the season, Crescent lake would be classified as a relatively clear and moderately productive (mesotrophic) lake.

9) Comparisons between Lay Monitor and FBG data indicate the volunteer monitor of Crescent Lake is doing an excellent job of measuring water quality at Site 6 Center. Alkalinity measurements collected by the volunteer monitor are slightly higher than readings taken by the FBG.

WENTWORTH LAKE

1992 NON-TECHNICAL SUMMARY

Monitoring was undertaken at Wentworth Lake by the volunteer monitors from June 3 to September 18. A more in-depth analysis of Wentworth Lake was conducted by the FBG on August 3.

1) Water transparency at Wentworth Lake was high, the sign of a clear and unproductive lake. The secchi disk was visible as far down as 9.4 meters (30.6 feet) and the transparency averages were 6.8 meters at Site 1 Fullers, 7.1 meters at Site 2 Triggs and 7.8 meters at Site 12 GovDeep. This indicates the deepwater sites on the lake contain little dissolved color and suspended matter such as algae and particulates. The average water clarity in 1992 is similar to the 1991 average.

2) Chlorophyll *a* concentrations for the surface waters of Wentworth Lake were low at all sampling stations in 1992. Concentrations in the mixed layer of water averaged 1.9 milligrams per cubic meter (1.9 mg m^{-3}, equivalent to about 1.9 parts chlorophyll per billion parts water) at Site 1 Fullers, 1.7 mg m^{-3} at Site 2 Triggs and 2.2 mg m^{-3} at Site 12 GovDeep. Generally, concentrations below 3 mg m^{-3} are common to less productive clear lakes and values above 7 mg m^{-3} are common in productive lakes. The 1992 average chlorophyll *a* concentrations are indicative of an unproductive, New Hampshire, lake.

3) Dissolved lakewater color levels for Wentworth Lake were low in 1992, 16.6 ptu (platinate color units), and less than the average levels of 26 ptu in other program lakes. Small increases in water color from the natural breakdown of plant materials in and around a lake are not considered to be detrimental to water quality. However, increased color can lower water transparency, and hence, change the public perception of water quality. Large amounts of dissolved color may occur naturally but also occur during deforestation and

development within the watershed. High color levels can actually mask the ability of the secchi disk transparency to predict chlorophyll levels.

4) Total phosphorus (nutrient) levels were low to moderate in Wentworth Lake when sampled by the volunteer monitors on June 9. Additional samples collected by the **FBG**, near the lake-bottom, were low when sampled on August 3 and exhibited no accumulation of phosphorous in the bottom waters. Tributary samples, collected by the volunteer monitors, reached moderate to high levels in the Tyler Brook, 21.8 ppb, and the Townsend Brook, 17.2 ppb, on August 14 and June 3, respectively. All other tributary samples remained below the concentration of 15 ppb commonly thought of as the boundary between less productive and more productive lakes.

5) The pH of the surface waters of the lake, measured by the **FBG** and volunteer monitors, remains within the optimum range for most aquatic organisms. The alkalinity of the lake, the lake's ability to buffer acid input, is low, and slightly less than the average of 6.3 units for **LLMP** program lakes. While the alkalinity level is low, it remains sufficient to buffer against fluctuations in pH caused by acid precipitation.

6) The specific conductivity of the deep sites on Wentworth Lake was low, between 38.6 and 43.4 micro-Siemans. High conductivity values can indicate the presence of septic leachate or deicing salt runoff.

7) Temperature profiles collected by the volunteer monitors disclosed the typical temperature stratification patterns for northern temperate lakes. With the upper mixed layer of water extending to 10 meters during the summer stratification period. The oxygen content of the bottom waters remained above 5 milligrams per liter (the minimum concentration required for the successful growth and reproduction of most coldwater fish) down to about 25 meters at Site 1 Fullers and to about 16 meters at Site 2 Triggs.

8) For all measurements considered and averaged for the season, Wentworth Lake would be classified as a clear and unproductive lake.

9) Comparisons between Lay Monitor and FBG data indicate the volunteer monitors of Wentworth Lake are doing an excellent job of measuring water quality at the deep sampling stations. Alkalinity measurements collected by the volunteer monitors are slightly higher than readings taken by the FBG.

COMMENTS AND RECOMMENDATIONS

1) We recommend that each Association, including the Wentworth Lake Association, continue to develop its data base on lake water quality through continuation of the long term monitoring program. The data base will provide information on the short and long-term cyclic variability that occurs in the lake and eventually will enable more reliable predictions of water quality trends.

2) We recommend continued phosphorous sampling in Crescent Lake late in the season to monitor the lakes reaction to a full seasons use. Additional sampling is recommended the week before and the week immediately following the Fourth of July to monitor the lakes response to increased lake usage during that period. If a boat is available, the **FBG** would like to schedule a team trip in early July to perform the typical sampling on Wentworth and Crescent Lakes as well undertake additional nutrient testing in Crescent Lake at that time.

3) With higher phosphorous (nutrient) concentrations occurring in the Ryfield and Townsend Brooks, we recommend the Wentworth Lake Association conduct a visual survey of the respective tributaries, noting any suspect activities occurring alongside the inlets which might be contributing to the elevated phosphorous levels as well as any other types of pollution (i.e. excessive sedimentation resulting from recent land clearing within the watershed).

INTRODUCTION

The New Hampshire Lakes Lay Monitoring Program

In this fifteenth year of operation, the **NH Lakes Lay Monitoring Program** has grown from a university class project on Chocorua Lake and pilot study on the Squam Lakes to a comprehensive state-wide program with over 500 volunteer monitors and more than 100 lakes participating. Originally developed to establish a data-base for determining long-term trends of lake water quality for science and management, the program has expanded by taking advantage of the many resources that citizen monitors can provide. The **NH LLMP** has an international reputation as a successful cooperative monitoring, education and research program. Current projects include: use of volunteer generated data for non-point pollution studies using high tech analysis system (Geographic Information Systems and Satellite Remote Sensing), intensive watershed monitoring for the development of lake nutrient budgets, and investigations of water quality and indicator organisms (food web analysis, fish condition, and stream invertebrates). The key ingredients responsible for the success of the program include innovative funding and cost reduction, assurance of credible data, practical sampling protocols and, most importantly, the interest and motivation of our volunteer monitors.

Importance of Long-term Monitoring

A major goal of a monitoring program is to identify any short or long-term changes in the water quality of the lake. Of major concern is the detection of cultural eutrophication: increases in the productivity of the lake, the amount of algae and plant growth, due to the addition of nutrients from human activities. Changes in the natural buffering capacity of the lakes in the program is also a topic of great concern, as New Hampshire receives large amounts of acid precipitation, yet most of our lakes contain little mineral content to neutralize this type of pollution.

For over a decade, data collected weekly from lakes participating in the New **Hampshire Lakes Lay Monitoring Program** have indicated there is quite a variation in water quality indicators through the open water season on the majority of lakes. Short-term differences may be due to variations in weather, lake use, or other chance events. Monthly sampling of a lake during a single summer provides some useful information, but there is a greater chance that important short-term events such as algal blooms or the lake response to storm run-off will be missed. These short-term fluctuations may be unrelated to the actual long-term trend of a lake or they may be indicative of the changing status or "health" of a lake.

To determine if a change in water quality is occurring, a lake must be sampled on a frequent basis over a substantial amount of time. A poorly designed sampling program may even mislead the investigator away from the actual trend: Consider the hypothetical lake in Figure 1. Sampling only once a year during August from 1982 to 1986 would produce a plot (Fig. 2) suggesting a decrease in eutrophication. The actual long-term trend of the lake, increasing eutrophy, can only be clearly discerned by sampling additional times a year for a ten year period (Fig. 1). Frequent monitoring carried out over the course of many summers can provide the information required to distinguish between short-term fluctuation ("noise") and long-term trends ("signal"). To that end, the lake must establish a long-term data base.

The number of seasons it takes to distinguish between the noise and the signal is not the same for each lake. Evaluation and interpretation of a long-term data base will indicate that the water quality of the lake has worsened, improved, or remained the same. In addition, different areas of a lake may show a different response. As more data is collected, prediction of current and future trends can be made. No matter what the outcome, this information is essential for the intelligent management of the lake.

There are also short-term uses for lay monitoring data. The examination of different stations in a lake can disclose the location of specific problems and corrective action can be initiated to handle the situation before it becomes more serious. On a lighter note, some associations post their weekly data for use in determining the best depths for finding fish!

It takes a considerable amount of effort as well as a deep concern for one's lake to be a lay monitor in the **NH Lakes Lay Monitoring Program.** Many times a monitor has to brave inclement weather or heavy boat traffic to collect samples. Sometimes it even may seem that one week's data is just the same as the next. Yet every sampling provides important information on the variability of the lake.

We are pleased with the interest and commitment of our Lay Monitors and are proud that their work is what makes the **NH LLMP** the most extensive, and we believe, the best volunteer program of its kind.

Purpose and Scope of This Study

This was the ninth year that monitoring of Wentworth and Crescent Lakes was undertaken by the **Freshwater Biology Group** and the Wentworth Lake Association. The program of sampling was designed to continue adding data to the long-term data base established. Sampling emphasis was placed on three open water deep stations in Wentworth Lake and one open water deep station in Crescent Lake while additional tributary sampling was undertaken at various locations around Wentworth Lake (See figure 3 for site locations).

The primary purpose of this report is to discuss results of the 1992 monitoring with emphasis on current conditions of Wentworth and Crescent Lakes including the extent of eutrophication and the lake's susceptibility to increasing acid precipitation. This information is part of a large data base of historical and more recent data compiled and entered onto computer files for New Hampshire lakes that include New Hampshire Fish

and Game surveys of the 1930's, the surveys by the New Hampshire Water Supply and Pollution Control Commission and the FBG surveys. Care must be taken when comparing current results with early studies. Many complications arise due to methodological differences of the various testing facilities and technological improvements in testing.

DISCUSSION OF LAKE MONITORING MEASUREMENTS

The section below details the important concepts involved for the various testing procedures used in the **New Hampshire Lakes Lay Monitoring Program.** Where appropriate, summary statistics of 1992 results from all participating lakes are included. Certain tests or sampling performed at the time of the optional **Freshwater Biology Group** field trip are indicated by an asterisk (*).

Thermal Stratification in the Deep Water Sites

Lakes in New Hampshire display distinct patterns of temperature stratification, that develop as the summer months progress, where a layer of warmer water (the **epilimnion**) overlies a deeper layer of cold water (**hypolimnion**). The layer that separates the two regions characterized by a sharp drop in temperature with depth is called the **thermocline** or **metalimnion.** Some shallow lakes may be continually mixed by wind action and will never stratify. Other lakes may only contain a developed epilimnion and metalimnion.

Wentworth Lake became stratified into three distinct layers, discussed above, as the season progressed while Crescent Lake is shallow and became only partially stratified when the weather was calm.

Water Transparency

Secchi Disk depth is a measure of the water transparency. The deeper the depth of secchi disk disappearance, the more transparent the lake water; light penetrates deeper if there is little dissolved and/or particulate matter (which includes both living and non-living particles) to absorb and scatter it.

In the shallow areas of many lakes, the secchi disk will hit bottom before it is able to disappear from view (what is referred to as a "Bottom Out" condition). Thus, Secchi disk measurements are generally taken over the deepest sites of a lake. Transparency values

of greater than 4 meters are typical of clear, less productive lakes. Values less than 2.5 meters are generally an indication of a very productive lake. In 1992 the average transparency for lakes participating in the NH LLMP was 5.6 meters with a range of 1.8 to 12.5 meters.

The average water clarity of Crescent Lake was low to moderate and averaged 4.9 meters (range: 4.0 to 5.6 meters) while the seasonal average water transparency was high in Wentworth Lake and averaged 6.8 meters (range: 5.7 to 8.2 meters) at Site 1 Fullers, 7.1 meters (range: 5.5 to 8.6 meters) at Site 2 Triggs and 7.8 meters (range: 5.8 to 9.4 meters) at Site 12 GovDeep. The seasonal average water clarity increased in both Wentworth and Crescent Lakes over the past two years. However, this is most likely attributable to the atypically low seasonal precipitation over that period and not a reflection of changing practices within the watershed. With 1993 off to a wet start, we expect increased nutrient loading through overland runoff which translates into higher chlorophyll *a* levels during the summer sampling season.

Chlorophyll a

The chlorophyll *a* concentration is a measurement of the standing crop of phytoplankton and is often used to classify lakes into categories of productivity called trophic states. **Eutrophic** lakes are highly productive with large concentrations of algae and aquatic plants due to nutrient enrichment. Characteristics include accumulated organic matter in the lake basin and lower dissolved oxygen in the bottom waters. Summer chlorophyll *a* concentrations average above 7 mg m^{-3} (7 milligrams per cubic meter; 7 parts per billion). **Oligotrophic** lakes have low productivity and low nutrient levels and average summer chlorophyll *a* concentrations are generally less than 3 mg m^{-3}. These lakes generally have cleaner bottoms and high dissolved oxygen levels throughout. **Mesotrophic** lakes are intermediate in productivity with concentrations of chlorophyll *a*

generally between 3 mg m^{-3} and 7 mg m^{-3}. In 1992 the average chlorophyll for lakes participating in the NH LLMP was 2.8 mg m^{-3} with a range of 0.4 to 18.5 mg m^{-3}.

With the below average precipitation in 1992, limiting nutrient runoff into the lake, chlorophyll *a* concentrations were again low, as they were in 1991. The seasonal average chlorophyll *a* concentration was 3.0 mg m^{-3} (range: 1.9 to 4.0 mg m^{-3}) in Crescent Lake, Site 6 Center, while the seasonal average chlorophyll *a* concentrations for Wentworth Lake were 1.9 mg m^{-3} (range: 1.4 to 3.1 mg m^{-3}) at Site 1 Fullers, 1.7 mg m^{-3} (range: 0.6 to 2.4 mg m^{-3}) at Site 2 Triggs and 2.2 mg m^{-3} (range: 1.5 to 4.7 mg m^{-3}) at Site 12 GovDeep.

Testing is sometimes done to check for **metalimnetic algal populations**, algae that layer out at the thermocline and generally go undetected if only epilimnetic (point or integrated) sampling is undertaken. Chlorophyll concentrations of a water sample collected in the thermocline is compared to the integrated epilimnetic sample. Greater chlorophyll levels of the point sample, in conjunction with microscopic examination of the samples (see Phytoplankton section below), confirm the presence of such a population of algae. These populations should be monitored as they may be an indication of increased nutrient loading into the lake.

Mid-lake chlorophyll *a* samples revealed no such algal population occurring in Wentworth Lake or Crescent Lake at the time of FBG sampling.

Dissolved Color

The dissolved color of lakes is generally due to dissolved organic matter from **humic substances**, which are naturally-occurring polyphenolic compounds leached from decayed vegetation. Highly colored or "stained" lakes have a "tea" color. Such substances generally do not threaten water quality except as they diminish sunlight penetration into deep waters. Increases in dissolved watercolor can be an indication of increased

development within the watershed as many land clearing activities (construction, deforestation, and the resulting increased run-off) add additional organic material to lakes. Natural fluctuations of dissolved color occur when storm events increase drainage from wetlands areas within the watershed. As suspended sediment is a difficult and expensive test to undertake, both dissolved color and chlorophyll information is important when interpreting the secchi disk transparency.

Dissolved color is measured on a comparative scale that uses standard chloroplatinate dyes and is designated as a color unit or ptu. Lakes with color below 10 ptu are very clear, 10 to 20 ptu are slightly colored, 20 to 40 ptu are lightly tea colored, 40 to 80 ptu are tea colored and greater than 80 ptu indicates highly colored waters. Generally the majority of New Hampshire lakes have color between 20 to 30 ptu.

Total Phosphorus

Of the two "nutrients" most important to the growth of aquatic plants, nitrogen and phosphorus, it is generally observed that phosphorus is the more limiting to plant growth, and therefore the more important to monitor and control. Phosphorus is generally present in lower concentrations, and its sources arise primarily through human related activity in a watershed. Nitrogen can be fixed from the atmosphere by many bloom-forming blue-green bacteria, and thus it is difficult to control. The total phosphorus includes all dissolved phosphorus as well as phosphorus contained in or adhered to suspended particulates such as sediment and plankton. As little as 15 parts per billion of phosphorus in a lake can cause an algal bloom.

Generally, in the more pristine lakes, phosphorus values are higher after spring melt when the lake receives the majority of runoff from its surrounding watershed. The nutrient is used by the algae and plants which in turn die and sink to the lake bottom causing phosphorus to decrease as the summer progresses. Lakes with nutrient loading

from human activities and sources (Agriculture, Sediment Erosion, Septic Systems, etc) will show greater concentrations of nutrients as the summer progresses or after major storm events. Circulation of nutrients from the bottom waters of more productive lakes in late fall can result in algal blooms.

Total Phosphorous levels were low to moderate in the surface waters of Wentworth and Crescent Lakes with a range of 4.0 to 10.4 ppb. Phosphorous samples collected by the FBG in early August revealed no accumulation of phosphorous in the bottom waters at that time. Additional samples collected at the inlets of Wentworth Lake were low to moderate with the exception of the Tyler Brook (21.8 ppb) and the Townsend Brook (17.2 ppb) which exceeded the concentration of 15 ppb on the August 14 and June 7 sampling dates, respectively (See figure 3 for site locations).

pH *

The pH is a way of expressing the acidic level of lake water, and is generally measured with an electrical probe sensitive to hydrogen ion activity. The pH scale has a range of 1 (very acidic) to 14 (very "basic" or alkaline) and is logarithmic (ie: changes in 1 pH unit reflect a ten times difference in hydrogen ion concentration). Most aquatic organisms tolerate a limited range of pH and most fish species require a pH of 5.5 or higher for successful growth and reproduction.

The pH of both Wentworth and Crescent Lakes was sufficient to support most aquatic life and ranged from 6.5 tp 6.7 units on August 3, 1992.

Alkalinity

Alkalinity is a measure of the buffering capacity of the lake water. The higher the value the more acid that can be neutralized. Typically lakes in New Hampshire have low alkalinities due to the absence of carbonates and other natural buffering minerals in the bedrock and soils of lake watersheds.

Decreasing alkalinity over a period of a few years can have serious effects on the lake ecosystem. In a study on an experimental acidified lake in Canada by Schindler, gradual lowering of the pH from 6.8 to 5.0 in an 8-year period resulted in the disappearance of some aquatic species, an increase in nuisance species of algae and a decline in the condition and reproduction rate of fish. During the first year of Schindler's study the pH remained unchanged while the alkalinity declined to 20 percent of the pre-treatment value. The decline in alkalinity was sufficient to trigger the disappearance of zooplankton species, which in turn caused a decline in the "condition" of fish species that fed on the zooplankton.

The analysis of alkalinity employed by the **Freshwater Biology Group** includes use of a dilute titrant allowing an order of magnitude greater sensitivity and precision than the standard method. Two endpoints are recorded during each analysis. The first endpoint (grey color of dye; pH endpoint of 5.1) approximates low level alkalinity values, while the second endpoint (pink dye color; pH endpoint of 4.6) approximates the alkalinity values recorded historically, such as NH Fish and Game data, with the methyl-orange endpoint method.

The average alkalinity of lakes throughout New Hampshire is low, approximately 9 mg per liter (calcium carbonate alkalinity), while the average alkalinity of the lakes studied by the **Freshwater Biology Group** in the **NH LLMP** is approximately 6.3 mg per liter. When alkalinity falls below 2 mg per liter the pH of waters can greatly fluctuate. Alkalinity levels are most critical in the spring when acid loadings from snowmelt and run-off are high, and many aquatic species are in their early, and most susceptible, stages of their life cycle.

The alkalinity of both Wentworth Lake (5.5 mg/L) and Crescent Lake (6.5 mg/L) remains low, but sufficient to buffer against wide fluctuations in pH caused by acid precipitation.

Specific Conductivity *

The specific conductance of a water sample indicates concentrations of dissolved salts. Leaking septic systems and deicing salt runoff from highways can cause high conductivity values. Fertilizers and other pollutants can also increase the conductivity of the water. Conductivity is measured in micromhos (the opposite of the measurement of resistance ohms) per centimeter, more commonly referred to as micro-Siemans.

The specific conductance of both Wentworth and Crescent Lakes remained low with a range of 38.6 to 43.4 micro-Siemans in the former, while the latter exhibited a range of 43.1 to 44.6 micro-Siemans.

Dissolved Oxygen and Free Carbon Dioxide *

Oxygen is an essential component for the survival of aquatic life. Submergent plants and algae take in free carbon dioxide and create oxygen through **photosynthesis** by day. **Respiration** by both animals and plants uses up oxygen continually and creates **carbon dioxide**. Dissolved oxygen profiles determine the extent of declining oxygen concentrations in the lower waters. High carbon dioxide values are indicative of low oxygen conditions and accumulating organic matter. For both gases, as the temperature of the water decreases, more gas can be dissolved in the water.

The typical pattern of clear, unproductive lakes is a slight decline in hypolimnetic oxygen as the summer progresses. Oxygen in the lower waters is important for maintaining a fit, reproducing, cold water fishery. Trout and salmon generally require oxygen concentrations above 5 mg per liter (parts per million) in the cool deep waters. On the other hand, carp and catfish can survive very low oxygen conditions. Oxygen above the lake bottom is important in limiting the release of nutrients from the sediments and minimizing the collection of undecomposed organic matter.

Bacteria, fungi and other **decomposers** in the bottom waters break down organic matter originating from the watershed or generated by the lake. This process uses up oxygen and produces carbon dioxide. In lakes where organic matter accumulation is high, oxygen depletion can occur. In highly stratified eutrophic lakes the entire hypolimnion can remain unoxygenated or **anaerobic** until fall mixing occurs.

The oxygen peaks occurring at surface and mid-lake depths during the day are quite common in many lakes. These characteristic **heterograde oxygen curves** are the result of the large amounts of oxygen, the by-product of photosynthesis, collecting in regions of high algal concentrations. If the peak occurs in the thermocline of the lake, metalimnetic algal populations (discussed above) may be present.

The dissolved oxygen concentrations remained high in both Wentworth and Crescent Lakes on the August 3 sampling date and well above the minimal levels required for the successful growth and reproduction for most warm-water fish species.

Underwater Light *

Underwater light available to photosynthetic organisms is measured with an **underwater photometer** which is much like the light meter of a camera (only waterproofed !). The **photic zone** of a lake is the volume of water capable of supporting photosynthesis. It is generally considered to be delineated by the water's surface and the level where light is reduced, by the absorption and scattering properties of the lake water, to one percent of the surface intensity. The one percent depth is sometimes termed the **compensation depth**. Knowledge of light penetration is important when considering lake productivity and in studies of submerged vegetation. Discontinuity (abrupt changes in the slope) of the profiles could be due to metalimnetic layering of algae or other particulates (discussed above). The underwater photometer allows the investigator to measure light at depths below the Secchi disk depth to supplement the transparency information.

Light profiles collected by the FBG on August 3 revealed the photic zone of Crescent Lake extended to the lake-bottom while the photic zone of Wentworth Lake extended to about 9.9 meters. That is to say, there is enough light to support plant growth over the entire lake-bottom of Crescent Lake while plants can only grow down to about 9.9 meters in Wentworth Lake.

Indicator Bacteria *

Coliform bacteria in water indicate the possibility of fecal contamination. Although they are usually considered harmless to humans, they are much easier to test for than harmful pathogenic enteric bacteria (**Salmonella, Shigella** etc.) and viruses that may be present in fecal material. **Total coliform** includes all coliform bacteria which arise from the gut of animals or from vegetative materials. **Fecal coliform** are those specific organisms that inhabit the gut of warm blooded animals. Another indicator organism **Fecal streptococcus** (sometimes referred to as **enterococcus**) also can be monitored. The ratio of fecal coliform to fecal strep may be useful in suggesting the type of animal source responsible for the contamination. Desirable levels for a Class A water body is less than 50 total coliform organisms per 100 milliliters. If the coliform level rises above 150 organisms per 100ml swimming should be prohibited.

Ducks and geese are often a common cause of high concentrations of coliform at specific lake sites. While waterfowl are important components to the natural and aesthetic qualities of lakes that we all enjoy, it is poor management practice to encourage these birds by feeding them. The lake and surrounding area provides enough healthy and natural food for the birds and feeding them stale bread or crackers does nothing more than import additional nutrients into the lake and allows for increased plant growth. As birds also are a host to the parasite that causes "swimmers itch" waterfowl roosting areas offer a greater

chance for infestation to occur. Thus while leaving offerings for our feathered friends is enticing, the results can prove to be detrimental to the lake system and to human health.

Phytoplankton *

The planktonic community includes microbial organisms that represent diverse life forms, containing photosynthetic as well as non-photosynthetic types, and including bacteria, algae, crustaceans and insect larvae (the zooplankton are discussed below in a separate section). Because planktonic algae or "phytoplankton" tend to undergo rapid seasonal cycles on a time scale of days and weeks, the levels of populations found should be considered to be most representative of the time of collection and not necessarily of other times during the ice-free season, especially the early spring and late fall periods.

The composition and concentration of phytoplankton can be indicative of the trophic status of a lake. Seasonal patterns do occur and must be considered. For example **diatoms**, tend to be most abundant in April-June and October-November, in the surface or epilimnetic layers of New Hampshire lakes. As the summer progresses, the dominant types might shift to **green algae** or **golden algae**. By late season **Blue-green bacteria** generally dominate. In nutrient rich lakes, nuisance green algae and/or bluegreen bacteria might dominate continually. After fall mixing diatoms might again be found to bloom.

The phytoplankton community of Wentworth Lake demonstrated a low density and a moderate to high diversity which is generally considered indicative of healthy lake conditions. Quantitative analysis of phytoplankton samples from Sites 1 Fullers and 2 Triggs indicate the dominant algae at the time of FBG sampling were small bluegreen bacteria filaments (trichomes), however, they were not of the nuisance variety. Crescent Lake demonstrated an increase in the concentration of bluegreen filaments which is indicative of a higher level of productivity commonly observed in shallow water bodies

such as Crescent Lake. However, the sample was not well preserved and thus only a qualitative analysis of this sample was possible.

Zooplankton *

There are three groups of zooplankton that are generally prevalent in lakes: the **protozoa, rotifers** and **crustaceans**. Most research has been devoted to the last two groups although protozoa may be found in substantial amounts. Of the rotifers and the crustaceans, time and budgetary constraints usually make it necessary to sample only the larger zooplankton (macrozooplankton; larger than 80 or 150 microns; 1 million microns make up a meter). Thus, zooplankton analysis is generally restricted only to the larger crustaceans. Crustacean zooplankton are very sensitive to pollutants and are commonly used to indicate the presence of toxic substances in water. The crustaceans can be divided into two groups, the **cladocerans** (which include the "water fleas") and the **copepods**.

Macrozooplankton are an important component in the lake system. The filter feeding of the herbivorous ("grazing") species may control the population size of selected species of phytoplankton. The larger zooplankton can be an important food source for juvenile and adult planktivorous fish. All zooplankton play a part in the recycling of nutrients within the lake.

As discussed above for phytoplankton, zooplankton undergo seasonal population cycles and the results discussed below are most representative of the collection dates and not necessarily of other times during the ice-free season, especially during the early spring and late fall.

The Macrozooplankton community of Wentworth Lake demonstrated a low to moderate density and high diversity, as did the phytoplankton assemblage. Site 1 Fullers was dominated by the cladoceran, *Holopedium*, while Site 2 Triggs was dominated by the *Cyclopoid* copepods which are commonly the most abundant form in the Macrozooplankton

community. The Macrozooplankton density of Crescent Lake was considerably higher than that in Wentworth Lake and reached a concentration of 26.6 animals per liter of water. The dominant zooplanktonic form was the cladoceran, *Diaphanosoma*, which tends to dominate in bacteria rich environments, as these organisms are selectively herbivorous upon smaller particles. As with the phytoplankton assemblage of Crescent Lake, the zooplankton population is indicative of a higher level of productivity than that observed in Wentworth Lake.

Fish Condition

As with the plankton discussed above, the health of the fish species of a lake will be indicative of the overall water quality. Condition is determined by comparing the length of the fish to its weight. As would be expected, the heavier the fish for its length, the better its condition will be. By also examining a scale collected from the fish under a microscope, the approximate age and growth history can also be determined.

REFERENCES

American Public Health Association.(APHA) 1985. Standard Methods for the Examination of Water and Wastewater 16th edition. APHA, AWWA, WPCF.

Baker, A.L. 1973. Microstratification of phytoplankton in selected Minnesota lakes. Ph. D. thesis, University of Minnesota.

Carlson, R.E. 1977. A trophic state index for lakes. Limnol. Oceanogr. 22:361-379.

Edmondson, W.T. 1937. Food conditions in some New Hampshire lakes. In: Biological survey of the Androscoggin, Saco and coastal watersheds. (Report of E.E. Hoover.) New Hampshire Fish and Game Commission, Concord, New Hampshire.

Forsberg, C. and S.O. Ryding. 1980. Eutrophication parameters and trophic state indices in 30 Swedish waste-water receiving lakes. Arch. Hydrobiol. 89:189-207

Gallup, D.N. 1969. Zooplankton distributions and zooplankton-phytoplankton relationships in a mesotrophic lake. Ph.D. Thesis, University of New Hampshire.

Haney, J.F. and D.J. Hall. 1973. Sugar-coated Daphnia: a preservation technique for Cladocera. Limnol. Oceanogr. 18:331-333.

Hoover, E.E. 1936. Preliminary biological survey of some New Hampshire lakes. Survey report no. 1. New Hampshire Fish and Game Department, Concord, New Hampshire.

Hoover, E.E. 1937. Biological survey of the Androscoggin, Saco, and coastal watersheds. Survey report no. 2. New Hampshire Fish and Game Department, Concord, New Hampshire.

Hoover, E.E. 1938. Biological Survey of the Merrimack watershed. Survey report no. 3. New Hampshire Fish and Game Department, Concord, New Hampshire.

Hutchinson, G.E. 1967. A treatise on limnology, vol. 2. John Wiley and Sons, New York.

Lind, O.T. 1979. Handbook of common methods in limnology. C.V. Mosby, St. Louis.

Lorenzen, M.W. 1980. Use of chlorophyll-Secchi disk relationships. Limnol. Oceanogr. 25:371-372.

New Hampshire Water Supply and Pollution Control Commission. 1981. Classification and priority listing of New Hampshire lakes. Staff report no. 121. Concord, New Hampshire.

Newell, A.E. 1977. Biological survey of the lakes and ponds in Sullivan, Merrimack, Belknap and Strafford Counties. Survey report no. 8b. New Hampshire Fish and Game Department, Concord, New Hampshire.

Schindler, D.W., et al. 1985. Long-term ecosystem stress: Effects of years of experimental acidification on a small lake. Science. 228:1395-1400.

Schloss, J.A., A.L. Baker and J.F. Haney. 1989. Over a decade of citizen volunteer monitoring in New Hampshire: The New Hampshire Lakes Lay Monitoring Program. Lake and Reservoir Management.

Sprules, W.G. 1980. Zoogeographic patterns in size structure of zooplankton communities with possible applications to lake ecosystem modeling and management. in W.C. Kerfoot ed. Evolution and Ecology of Zooplankton Communities. University Press of New England. Dartmouth. pp642-656.

Uttermohl, H. 1958. Improvements in the quantitative methods of phytoplankton study. Mitt. int. Ver. Limnol. 9:1-25.

U.S. Environmental Protection Agency. 1979. A manual of methods for chemical analysis of water and wastes. Office of Technology Transfer, Cincinnati. PA-600/4-79-020.

Vollenweider, R.A. 1969. A manual on methods for measuring primary productivity in aquatic environments. International Biological Programme. Blackwell Scientific Publications, Oxford.

Warfel, H.E. 1939. Biological survey of the Connecticut Watershed. Survey Report 4. N.H. Fish and Game. Concord, New Hampshire.

Wetzel, R.G. 1983. Limnology. Saunders College Publishing, Philadelphia.

Wetzel, R.G. and G.E. Likens. 1979. Limnological Analyses. W.B. Saunders Co. Philadelphia.

REPORT FIGURES



.Figure 1. The upper graph depicts weekly chlorophyll concentrations of a model lake measured weekly during ice-free conditions. The long-term trend is that of increased eutrophication (lake has become "greener"). Diamonds below the curve represent late summer (August) dates the data set was subsampled to create Figure 2.



Figure 2. The lower graph depicts late summer chlorophyll data of the model lake in Figure 1. Note how limited sampling over a five year period suggests a much different trend, that of decreasing eutrophy. Thus, limited sampling can mislead the investigator of long-term trends.

Figure 3. Location of 1992 sampling stations for Wentworth Lake, Wolfeboro, New Hampshire.

11 Fernald Bk
3 Willey Bk
10 Clay Pit Bk
4 Tyler Bk
5B Frost Bk
5 Ryfield Bk
2 Triggs
12 GovDeep
1 Fullers
9 Warren
8 Townsend Bk
7 Heath Bk

Figure 4. Crescent Lake, Site 6 Center. Seasonal trends for Secchi Disk Depth (water transparency), 1992. Dotted lines on the plot border the ranges common to oligotrophic, mesotrophic and eutrophic lakes. The dashed horizontal line on the plot indicates the maximum depth at the site.

Figure 5. Crescent Lake, 1992. Seasonal trends for chlorophyll *a* concentration of Lay Monitor Site 6 Center. Chlorophyll *a* concentrations in parts per billion (ppb) of chlorophyll *a*. The dotted horizontal lines distinguish the ranges common to oligotrophic, mesotrophic and eutrophic lakes.

Figure 6. Crescent Lake, 1992. Seasonal trends for dissolved color concentration of Lay Monitor Site 6 Center. Color expressed as platinum-cobalt units (ptu). The dotted line represents the 1992 average dissolved color level for participating **LLMP** lakes.

CRESCENT LAKE — 6 CENTER

SECCHI DISK TRANSPARENCY 1992



CRESCENT LAKE

CHLOROPHYL CONCENTRATION 1992



CRESCENT LAKE

DISSOLVED COLOR CONCENTRATION 1992



Figure 7. Wentworth Lake, Site 1 Fullers. Seasonal trends for Secchi Disk Depth (water transparency), 1992. Dotted lines on the plot border the ranges common to oligotrophic, mesotrophic and eutrophic lakes.

Figure 8. Wentworth Lake, 1992. Seasonal trends for chlorophyll *a* concentration of Lay Monitor Site 1 Fullers. Chlorophyll *a* concentrations in parts per billion (ppb) of chlorophyll *a*. The dotted horizontal lines distinguish the ranges common to oligotrophic, mesotrophic and eutrophic lakes.

Figure 9. Wentworth Lake, 1992. Seasonal trends for dissolved color concentration of Lay Monitor Site 1 Fullers. Color expressed as platinum-cobalt units (ptu). The dotted line represents the 1992 average dissolved color level for participating **LLMP** lakes.

LAKE WENTWORTH — 1 FULLERS

SECCHI DISK TRANSPARENCY 1992



LAKE WENTWORTH

CHLOROPHYL CONCENTRATION 1992



LAKE WENTWORTH

DISSOLVED COLOR CONCENTRATION 1992



Figure 10. Wentworth Lake, Site 2 Triggs. Seasonal trends for Secchi Disk Depth (water transparency), 1992. Dotted lines on the plot border the ranges common to oligotrophic, mesotrophic and eutrophic lakes.

Figure 11. Wentworth Lake, 1992. Seasonal trends for chlorophyll *a* concentration of Lay Monitor Site 2 Triggs. Chlorophyll *a* concentrations in parts per billion (ppb) of chlorophyll *a*. The dotted horizontal lines distinguish the ranges common to oligotrophic, mesotrophic and eutrophic lakes.

Figure 12. Wentworth Lake, 1992. Seasonal trends for dissolved color concentration of Lay Monitor Site 2 Triggs. Color expressed as platinum-cobalt units (ptu). The dotted line represents the 1992 average dissolved color level for participating **LLMP** lakes.

LAKE WENTWORTH — 2 TRIGGS

SECCHI DISK TRANSPARENCY 1992



LAKE WENTWORTH

CHLOROPHYL CONCENTRATION 1992



\+ 2 TRIGGS

LAKE WENTWORTH

DISSOLVED COLOR CONCENTRATION 1992



\+ 2 TRIGGS

Figure 13. Wentworth Lake, Site 12 GovDeep. Seasonal trends for Secchi Disk Depth (water transparency), 1992. Dotted lines on the plot border the ranges common to oligotrophic, mesotrophic and eutrophic lakes.

Figure 14. Wentworth Lake, 1992. Seasonal trends for chlorophyll *a* concentration of Lay Monitor Site 12 GovDeep. Chlorophyll *a* concentrations in parts per billion (ppb) of chlorophyll *a*. The dotted horizontal lines distinguish the ranges common to oligotrophic, mesotrophic and eutrophic lakes.

Figure 15. Wentworth Lake, 1992. Seasonal trends for dissolved color concentration of Lay Monitor Site 12 GovDeep. Color expressed as platinum-cobalt units (ptu). The dotted line represents the 1992 average dissolved color level for participating **LLMP** lakes.

LAKE WENTWORTH — 12 GOVDEEP

SECCHI DISK TRANSPARENCY 1992



LAKE WENTWORTH

CHLOROPHYL CONCENTRATION 1992



LAKE WENTWORTH

DISSOLVED COLOR CONCENTRATION 1992



Figure 16. Lake Wentworth, 1992. Seasonal trends for chlorophyll *a* concentration of Lay Monitor Sites 1 Fullers (squares), 2 Triggs (crosses) and 12 GovDeep (diamonds). Chlorophyll *a* concentrations in parts per billion (ppb) of chlorophyll *a*. The dotted lines on the plot border the ranges common to oligotrophic, mesotrophic and eutrophic lakes.

Figure 17. Lake Wentworth, 1992. Seasonal trends for dissolved color concentration of Lay Monitor Sites 1 Fullers (squares), 2 Triggs (crosses) and 12 GovDeep (diamonds). Color expressed as platinum-cobalt units (ptu). The dotted horizontal line represents the average dissolved color concentration for participating **LLMP** lakes.

LAKE WENTWORTH

CHLOROPHYL CONCENTRATION 1992



LAKE WENTWORTH

DISSOLVED COLOR CONCENTRATION 1992



Figure 18. Profiles of temperature (TEMP) and dissolved oxygen (DO) taken on August 3, 1992 at Wentworth Lake (A) Site 1 Fullers and (B) Site 2 GovDeep. Units of measurement are as indicated. Oxygen and temperature were measured at one-half meter intervals.

TEMPERATURE - OXYGEN PROFILE
WENTWORTH - SITE 1 FULLERS
AUGUST 3, 1992



TEMPERATURE - OXYGEN PROFILE
WENTWORTH - SITE 2 TRIGGS
AUGUST 3, 1992



Figure 19. Profile of temperature (TEMP) and dissolved oxygen (DO) taken on August 3, 1992 at Crescent Lake, Site 6 Center. Units of measurement are as indicated. Oxygen and temperature were measured at one-half meter intervals.

TEMPERATURE - OXYGEN PROFILE
CRESENT LAKE - SITE 6 CENTER
AUGUST 3, 1992



Figure 20. Pie diagrams of Phytoplankton Abundance collected in Lake Wentworth, Sites (A) 1 Fullers and (B) 2 Triggs on August 3, 1992. Phytoplankton abundance is presented as relative percent by algal group.

WENTWORTH LAKE

SITE 1 FULLERS

DEPTH 0-8.5 meters



SITE 2 TRIGGS

DEPTH 0-9.0 meters



RELATIVE PERCENT BY ALGAL GROUP

Figure 21. Pie diagrams of Macro-Zooplankton Diversity collected in Wentworth Lake Sites (A) 1 Fullers and (B) 2 Triggs on August 3, 1992. Macro-Zooplankton abundances are presented as number of animals per liter of lakewater.

LAKE WENTWORTH

1 FULLERS
MACROZOOPLANKTON DATA 0-24m
8-3-92



2 TRIGGS
MACROZOOPLANKTON DATA 0-15m
8-3-92



NUMBERS SHOWN ARE # OF ANIMALS PER LITER

Figure 22. Pie diagram of Macro-Zooplankton Diversity collected in Crescent Lake, Site 6 Center, on August 3, 1992. The Macro-Zooplankton abundance is presented as number of animals per liter of lakewater.

CRESCENT LAKE - SITE 6 CENTER
MACROZOOPLANKTON DATA 0-4.5m
8-3-92



NUMBERS SHOWN ARE # OF ANIMALS PER LITER

Figure 23. Comparison of 1992 Crescent Lake Chlorophyll *a* data to historical data (1984-1991). The patterns on the bars distinguish the minimum, mean and maximum values while the length of the bars represents the total range of values. The Chlorophyll *a* concentration is measured in parts per billion (ppb) which is equivalent to milligrams per cubic meter. The higher the chlorophyll *a* levels the "greener" the water (more algal growth).

Figure 24. Comparison of 1992 Crescent Lake secchi disk transparency data to historical data (1984-1991). The patterns on the bars distinguish the minimum, mean and maximum values while the length of the bars represents the total range of values. Secchi disk readings are taken to the nearest 0.1 meter. The deeper the secchi disk depth the clearer the water.

COMPARISON: 1992 TO HISTORICAL DATA
CRESCENT LAKE CHLOROPHYLL a
LAY MONITOR DATA



The higher number = more algae

COMPARISON: 1992 TO HISTORICAL DATA
CRESCENT LAKE WATER CLARITY
LAY MONITOR DATA



The higher number = clearer water

Figure 25. Comparison of 1992 Wentworth Lake Chlorophyll *a* data to historical data (1984-1991). The patterns on the bars distinguish the minimum, mean and maximum values while the length of the bars represents the total range of values. The Chlorophyll *a* concentration is measured in parts per billion (ppb) which is equivalent to milligrams per cubic meter. The higher the chlorophyll *a* levels the "greener" the water (more algal growth).

Figure 26. Comparison of 1992 Wentworth Lake secchi disk transparency data to historical data (1984-1991). The patterns on the bars distinguish the minimum, mean and maximum values while the length of the bars represents the total range of values. Secchi disk readings are taken to the nearest 0.1 meter. The deeper the secchi disk depth the clearer the water.

COMPARISON: 1992 TO HISTORICAL DATA
WENTWORTH LAKE CHLOROPHYLL *a*
LAY MONITOR DATA



The higher number = more algae

COMPARISON: 1992 TO HISTORICAL DATA
WENTWORTH LAKE WATER CLARITY
LAY MONITOR DATA



The higher number = clearer water

Lake Wentworth Data on file as of 12/29/1992

Lakes Lay Monitoring Program, U.N.H.

[Lay Monitor Data]

Lake Wentworth, NH
-- subset of trophic indicators, all sites, 1992

1992 SUMMARY

Average transparency:	7.2	(1992:	43	values;	5.5	-	9.4	range)
Average chlorophyll:	1.9	(1992:	44	values;	0.6	-	4.7	range)
Average phosphorus:	6.8	(1992:	2	values;	4.0	-	9.7	range)
Average alk (gray):	5.5	(1992:	43	values;	4.9	-	5.9	range)
Average alk (pink):	6.1	(1992:	43	values;	5.7	-	6.5	range)
Average color, 440:	16.6	(1992:	42	values;	11.2	-	37.8	range)
Average Trib. phos:	5.3	(1992:	32	values;	1.1	-	21.8	range)

Site	Date	Trans-parency (m)	Chl a (ppb)	Total Phos (ppb)	Alk. (gray) ph 5.1	Alk. (pink) ph 4.6	Color Pt-Co units
1 Fullers	06/09/1992	6.2	1.8	4.0	4.9	6.1	---
1 Fullers	06/16/1992	5.7	2.2	---	5.4	6.3	14.6
1 Fullers	06/24/1992	6.2	2.0	---	5.3	5.9	14.6
1 Fullers	06/29/1992	6.2	1.9	---	5.2	6.1	12.0
1 Fullers	07/08/1992	6.4	1.5	---	5.4	6.2	12.9
1 Fullers	07/14/1992	7.1	3.1	---	---	---	13.7
1 Fullers	07/20/1992	6.6	1.9	---	5.4	6.1	18.0
1 Fullers	07/26/1992	6.9	1.7	---	5.6	6.1	15.5
1 Fullers	08/02/1992	6.7	1.6	---	5.6	6.1	18.9
1 Fullers	08/09/1992	6.9	2.1	---	5.5	6.0	15.5
1 Fullers	08/15/1992	7.4	2.1	---	5.3	5.8	25.8
1 Fullers	08/22/1992	---	1.4	---	5.4	5.9	31.8
1 Fullers	09/01/1992	7.7	1.8	---	5.3	5.8	16.3
1 Fullers	09/06/1992	7.4	2.2	---	5.5	5.9	12.9
1 Fullers	09/13/1992	8.2	1.4	---	5.5	6.0	17.2
2 Triggs	06/09/1992	6.7	1.6	9.7	5.2	6.2	---
2 Triggs	06/16/1992	5.5	2.1	---	5.2	6.2	15.5
2 Triggs	06/24/1992	6.0	2.0	---	5.0	5.8	13.7
2 Triggs	06/29/1992	6.4	1.6	---	5.3	6.5	14.6
2 Triggs	07/08/1992	6.6	1.6	---	5.7	6.5	13.7
2 Triggs	07/14/1992	7.0	2.4	---	5.6	6.2	15.5
2 Triggs	07/20/1992	6.9	1.9	---	5.4	6.0	28.3
2 Triggs	07/26/1992	7.2	1.8	---	5.4	5.9	37.8
2 Triggs	08/02/1992	7.3	1.8	---	5.4	6.0	16.3
2 Triggs	08/09/1992	7.2	1.7	---	5.4	5.9	18.9
2 Triggs	08/15/1992	7.6	1.9	---	5.5	6.0	15.5
2 Triggs	08/22/1992	7.5	0.6	---	5.3	5.9	12.9
2 Triggs	09/01/1992	7.5	1.6	---	5.4	6.0	23.2
2 Triggs	09/06/1992	8.2	1.6	---	5.4	5.9	12.0
2 Triggs	09/13/1992	8.6	1.4	---	5.7	6.3	13.7
3 Shore	03/04/1992	---	---	4.2	---	---	---

Lake Wentworth Data on file as of 12/29/1992

Site	Date	Trans-parency (m)	Chl a (ppb)	Total Phos (ppb)	Alk. (gray) ph 5.1	Alk. (pink) ph 4.6	Color Pt-Co units
3 Shore	06/07/1992	---	---	1.1	---	---	---
3 Shore	08/14/1992	---	---	1.3	---	---	---
3 Shore	09/12/1992	---	---	7.7	---	---	---
4 Shore	06/07/1992	---	---	1.8	---	---	---
4 Shore	08/14/1992	---	---	21.8	---	---	---
4 Shore	09/12/1992	---	---	5.1	---	---	---
4 Tyler	06/03/1992	---	---	13.2	---	---	---
5 Ryfield	06/03/1992	---	---	12.8	---	---	---
5 Shore	06/07/1992	---	---	5.1	---	---	---
5 Shore	08/14/1992	---	---	1.7	---	---	---
5 Shore	09/12/1992	---	---	3.3	---	---	---
7 Heath	06/03/1992	---	---	12.1	---	---	---
7 Shore	06/07/1992	---	---	2.0	---	---	---
7 Shore	08/14/1992	---	---	1.3	---	---	---
7 Shore	09/12/1992	---	---	2.6	---	---	---
8 Shore	06/07/1992	---	---	1.1	---	---	---
8 Shore	08/14/1992	---	---	6.0	---	---	---
8 Shore	09/12/1992	---	---	1.5	---	---	---
8 Townsnd	06/03/1992	---	---	17.2	---	---	---
9 Shore	06/07/1992	---	---	2.4	---	---	---
9 Shore	08/14/1992	---	---	3.3	---	---	---
9 Shore	09/12/1992	---	---	4.4	---	---	---
9 Warren	06/02/1992	---	---	11.9	---	---	---
10 Clay	06/03/1992	---	---	2.9	---	---	---
10 Shore	06/07/1992	---	---	1.8	---	---	---
10 Shore	08/14/1992	---	---	1.1	---	---	---
10 Shore	09/12/1992	---	---	8.8	---	---	---
11 Fernald	06/05/1992	---	---	2.0	---	---	---
11 Shore	06/07/1992	---	---	2.0	---	---	---
11 Shore	08/14/1992	---	---	4.4	---	---	---
11 Shore	09/12/1992	---	---	1.1	---	---	---
12 GovDeep	06/13/1992	7.0	2.0	---	5.5	5.7	13.7
12 Govdeep	06/20/1992	5.8	1.9	---	5.6	6.0	13.7
12 GovDeep	06/27/1992	6.2	1.9	---	5.9	6.3	17.2
12 GovDeep	07/05/1992	6.9	4.7	---	5.5	5.8	12.9
12 GovDeep	07/11/1992	7.4	1.7	---	5.8	6.3	16.3
12 GovDeep	07/24/1992	8.4	2.7	---	5.4	5.8	16.3
12 GovDeep	08/02/1992	7.3	1.9	---	5.3	5.8	12.9
12 GovDeep	08/07/1992	7.8	2.3	---	5.5	6.0	12.0
12 GovDeep	08/14/1992	7.6	2.5	---	5.7	6.0	12.0
12 GovDeep	08/21/1992	8.0	2.0	---	5.5	6.1	11.2
12 GovDeep	08/27/1992	8.9	1.8	---	5.7	6.3	20.6
12 GovDeep	09/05/1992	9.4	1.5	---	5.7	6.1	12.0
12 GovDeep	09/11/1992	9.4	2.0	---	5.7	6.2	15.5
12 GovDeep	09/18/1992	9.3	1.6	---	5.8	6.2	18.0

Lake Wentworth Data on file as of 12/29/1992

<< End of 1992 listing, 76 records >>

Lake Wentworth Data on file as of 12/29/1992

Lakes Lay Monitoring Program, U.N.H.

[Lay Monitor Data]

Crescent Lake, NH
-- subset of trophic indicators, all sites, 1992

1992 SUMMARY

Average transparency:	4.9	(1992:	13	values;	4.0	-	5.6	range)
Average chlorophyll:	3.0	(1992:	13	values;	1.9	-	4.0	range)
Average phosphorus:	10.4	(1992:	1	values;	10.4	-	10.4	range)
Average alk (gray):	6.5	(1992:	13	values;	6.0	-	6.9	range)
Average alk (pink):	7.3	(1992:	13	values;	6.9	-	8.0	range)
Average color, 440:	18.3	(1992:	13	values;	12.0	-	34.4	range)

Site	Date	Trans-parency (m)	Chl a (ppb)	Total Phos (ppb)	Alk. (gray) ph 5.1	Alk. (pink) ph 4.6	Color Pt-Co units
6 Center	06/16/1992	4.2	1.9	---	6.0	7.1	18.9
6 Center	06/23/1992	5.0	3.1	---	6.5	7.2	34.4
6 Center	06/30/1992	4.5	2.7	---	6.9	8.0	16.3
6 Center	07/08/1992	5.1	2.3	---	6.5	7.1	17.2
6 Center	07/16/1992	4.9	2.9	---	6.4	6.9	14.6
6 Center	07/21/1992	5.0	3.1	---	6.5	7.0	15.5
6 Center	07/28/1992	5.6	2.6	---	6.4	7.3	13.7
6 Center	08/11/1992	4.7	2.9	---	6.6	7.2	12.0
6 Center	08/19/1992	5.2	4.0	---	6.4	7.7	12.0
6 Center	08/25/1992	4.0	3.0	---	6.4	7.0	28.3
6 Center	09/01/1992	4.9	3.3	---	6.8	7.4	24.1
6 Center	09/08/1992	5.4	3.9	---	6.4	7.0	15.5
6 Center	09/12/1992	---	---	10.4	---	---	---
6 Center	09/16/1992	4.7	3.9	---	6.9	7.4	15.5

<< End of 1992 listing, 14 records >>





TYPICAL TEMPERATURE CONDITIONS : SUMMER

NEW HAMPSHIRE - DEEP LAKE



APPENDIX C

GLOSSARY OF LIMNOLOGICAL TERMS

Aerobe- Organisms requiring oxygen for life. All animals, most algae and some bacteria require oxygen for respiration.

Algae- See phytoplankton.

Alkalinity- Total concentration of bicarbonate and hydroxide ions (in most lakes). Alkalinity is often referred to as the acid neutralizing capacity (ANC) in the literatue and alkalinity concentrations are generally recorded as parts per million (ppm), milliequivalents per liter and milligrams per liter.

Anaerobe- Organisms not requiring oxygen for life. Some algae and many bacteria are able to respire or ferment without using oxygen.

Anoxic- A system lacking oxygen, therefore incapable of supporting the most common kind of biological respiration, or of supporting oxygen-demanding chemical reactions. The deeper waters of a lake may become anoxic if there are many organisms depleting oxygen via respiration, and there is little or no replenishment of oxygen from photosynthesis or from the atmosphere.

Benthic- Referring to the bottom sediments.

Bacterioplankton- Bacteria adapted to the "open water" or "planktonic" zone of lakes, adapted for many specialized habitats and include groups that can use the sun's energy (phytoplankton), some that can use the energy locked in sulfur or iron, and others that gain energy by decomposing dead material.

Bicarbonate- The most important ion (chemical) involved in the buffering system of New Hampshire lakes.

Buffering- The capacity of lakewater to absorb acid with a minimal change in the pH. In New Hampshire the chemical responsible for buffering is the bicarbonate ion. (See pH.)

Chloride- One of the components of salts dissolved in lakewater. Generally the most abundant ion in New Hampshire lakewater, it may be used as an indicator of raw sewage or of road salt.

Chlorophyll a- The main green pigment in plants. The concentration of chlorophyll *a* in lakewater is often used as an indicator of algal abundance.

Circulation- The period during spring and fall when the combination of low water temperature and wind cause the water column to mix freely over its entire depth.

Density- The weight per volume of a substance. The more dense an object, the heavier it feels. Low-density liquids will float on higher-density liquids.

Dimictic- The thermal pattern of lakes where the lake circulates, or mixes, twice a year. Other patterns such as polymictic (many periods of circulation per year) are uncommon in New Hampshire. (See also meromictic and holomictic).

Dystrophy- The lake trophic state in which the lakewater is highly stained with humic acids (reddish brown or yellow stain) and has low productivity. Chlorophyll *a* concentration may be low or high.

Epilimnion- The uppermost layer of water during periods of thermal stratification. (See lake diagram).

Eutrophy- The lake trophic state in which algal production is high. Associated with eutrophy is low Secchi disk depth, high chlorophyll *a*, and high total phosphorus. From an esthetic viewpoint these lakes are "bad" because water clarity is low, aquatic plants are often found in abundance, and cold-water fish such as trout and salmon are usually not present. A good aspect of eutrophic lakes is their high productivity in terms of warm-water fish such as bass, pickerel, and perch.

Free CO2- Carbon dioxide that is not combined chemically with lake water or any other substances. It is produced by respiration, and is used by plants and bacteria for photosynthesis.

Holomixis- The condition where the entire lake is free to circulate during periods of overturn. (See meromixis.)

Humic Acids-Dissolved organic compounds released from decomposition of plant leaves and stems. Humic acids are red, brown, or yellow in color and are present in nearly all lakes in New Hampshire. Humic acids are consumed only by fungi, and thus are relatively resistant to biological decomposition.

Hydrogen Ion- The "acid" ion, present in small amounts even in distilled water, but contributed to rain-water by atmospheric processes, to ground-water by soils, and to lakewater by biological organisms and sediments. The active component of "acid rain". See also "pH" the symbolic value inversely and exponentially related to the hydrogen ion.

Hypolimnion-The deepest layer of lakewater during periods of thermal stratification. (See lake diagram)

Lake- Any "inland" body of relatively "standing" water. Includes many synonyms such as ponds, tarns, loches, billabongs, bogs, marshes, etc.

Lake Morphology- The shape and size of a lake and its basin.

Littoral- The area of a lake shallow enough for submerged aquatic plants to grow.

Meromixis- The condition where the entire lake fails to circulate to its deepest points; caused by a high concentration of salt in the deeper waters, and by peculiar landscapes (small deep lakes surrounded by hills and/or forests. (Contrast holomixis.)

Mesotrophy- The lake trophic state intermediate between oligotrophy and eutrophy. Algal production is moderate, and chlorophyll *a*, Secchi disk depth, and total phosphorus are also moderate. These lakes are esthetically "fair" but not as good as oligotrophic lakes.

Metalimnion- The "middle" layer of the lake during periods of summer thermal stratification. Usually defined as the region where the water temperature changes at least one degree per meter depth. Also called the thermocline.

Mixis- Periods of lakewater mixing or circulation.

Mixotrophy- The lake condition where the water is highly stained with humic acids, but algal production and chlorophyll *a* values are also high.

Oligotrophy- The lake trophic state where algal production is low, Secchi disk depth is deep, and chlorophyll *a* and total phosphorus are low. Esthetically these lakes are the "best" because they are clear and have a minimum of algae and aquatic plants. Deep oligotrophic lakes can usually support cold-water fish such as lake trout and land-locked salmon.

Overturn- See circulation or mixis

pH- A measure of the hydrogen ion concentration of a liquid. For every decrease of 1 pH unit, the hydrogen ion concentration increases 10 times. Symbolically, the pH value is the "negative logarithm" of the hydrogen ion concentration. For example, a pH of 5 represents a hydrogen ion concentration of 10^{-5} molar. [Please thank the chemists for this lovely symbolism -- and ask them to explain it in lay terms!] In any event, the higher the pH value, the lower the hydrogen ion concentration. The range is 0 to 14, with 7 being neutral 1 denoting high acid condition and 14 denoting very basic condition.

Photosynthesis- The process by which plants convert the inorganic substances carbon dioxide and water into organic glucose (sugar) and oxygen using sunlight as the energy source. Glucose is an energy source for growth, reproduction, and maintenance of almost all life forms.

Phytoplankton- Microscopic algae which are suspended in the "open water" zone of lakes and ponds. A major source of food for zooplankton. Common examples include: diatoms, euglenoids, dinoflagellates, and many others. Usually included are the blue-green bacteria.

Parts per million- Also known as "ppm". This is a method of expressing the amount of one substance (solute) dissolved in another (solvent). For example, a solution with 10 ppm of oxygen has 10 pounds of oxygen for every 999,990 pounds (500 tons) of water. Domestic sewage usually contains from 2 to 10 ppm phosphorus.

Parts per billion- Also known as "ppb". This is only 1/1000 of ppm, therefore much less concentrated. As little as 1 ppb of phosphorus will sustain growth of algae. As little as 10 ppb phosphorus will cause algal blooms! Think of the ratio as 1 milligram (1/28000 of an ounce) of phosphorus in 25 barrels of water (55 gallon drums)! Or, 1 gallon of septic waste diluted into 10,000 gallons of lakewater. It adds up fast!

Plankton- Community of microorganisms that live suspended in the water column, not attached to the bottom sediments or aquatic plants. See also "bacterioplankton" (bacteria), "phytoplankton" (algae) and "zooplankton" (microcrustaceans and rotifers).

Saturated- When a solute (such as water) has dissolved all of a substance that it can. For example, if you add table salt to water, a point is reached where any additional salt fails to dissolve. The water is then said to be saturated with table salt. In lakewater, gaseous oxygen can dissolve, but eventually the water becomes saturated with oxygen if exposed sufficiently long to the atmosphere or another source of oxygen.

Specific Conductivity- A measure of the amount of salt present in lakewater. As the salt concentration increases, so does the specific conductivity (electrical conductivity).

Stratum- A layer or "blanket". Can be used to refer to one of the major layers of lakewater such as the epilimnion, or to any layers of organisms or chemicals that may be present in a lake.

Thermal Stratification- The process by which layers are built up in the lake due to heating by the sun and partial mixing by wind.

Thermocline- Region of temperature change. (See metalimnion.)

Total Phosphorus- A measure of the concentration of phosphorus in lakewater. Includes both free forms (dissolved), and chemically combined form (as in living tissue, or in dead but suspended organisms).

Trophic Status- A classification system placing lakes into similar groups according to their amount of algal production. (See Oligotrophy, Mesotrophy, Eutrophy, Mixotrophy, and Dystrophy for definitions of the major categories)

Z- A symbol used by limnologists as an abbreviation for depth.

Zooplankton- Microscopic animals in the planktonic community. Some are called "water fleas", but most are known by their scientific names. Scientific names include: *Daphnia*, *Cyclops*, *Bosmina*, and *Kellicottia*.