

## Silviculture

Art & science of establishing & tending trees & forests



Karen Bennett, [karen.bennett@unh.edu](mailto:karen.bennett@unh.edu)  
Extension Forestry Professor & Specialist  
Presented to NH Coverts, May 2016



University of  
New Hampshire  
Cooperative Extension

## Silviculture Actions Have Two Broad Outcomes

- Grow the trees that are already present
  - tending
- Start new trees
  - regenerating
- In practice, often accomplish both outcomes at once
- Most common actions- cut trees or leave trees

Harvesting is the most common tool for conducting silviculture



## Forest Management/ Forest Stewardship

Interaction of silviculture, ecology, landowner objectives, multiple resources, economics, marketing, regulation, societies' needs and a landowner's interests and time.

- Markets, plans, laws, harvesting, equipment, landowner, logger, forester, neighbors, trails, access


## Silviculture

is the set of site specific tools used in forest management

- weeding, thinning, pruning, improving, harvesting, regenerating, uneven-aged, even-aged, selection, shelterwood, clearcut

## Hallmarks of Good Forest Stewardship/ Management

- Considers multiple resources
- Based on landowner objectives
- Uses best available practices
- Practices based on a plan
- Looks long term
- Uses professionals
- Uses best available science- SILVICULTURE



Silviculture can be used to create and maintain the kind of forest meets landowner objectives



Can be single or multiple objectives




Grow big trees for beauty, wildlife, and timber for money to send the kids to college.





Create standing snags and down woody material



Create temporary wildlife openings












## Shade Tolerance

Tolerance is the ability of a tree to grow satisfactorily in the shade of another tree.

As a stand succeeds tolerant species replace intolerant species.




## Tolerant vs. Intolerant

- Intolerant to shade: sun-requiring
  - tends not to reproduce under self
  - “sun-loving”
  - tend to be light seeded, wind-dispersed
  - early successional species
- Tolerant to shade: shade-adapted
  - reproduce under self
  - tend to be heavier seeded and moved by gravity or animals
  - later successional
- Intermediate

Softwoods		Hardwoods	
<b>Extremely Tolerant</b>			
balsam fir		American beech	
eastern hemlock		sugar maple	
<b>Tolerant</b>			
red spruce	white spruce		red maple
	northern white cedar		
<b>Intermediate</b>			
eastern white pine		white ash	red oak
		yellow birch	
<b>Intolerant</b>			
red pine			paper birch
<b>Extremely Intolerant</b>			
			aspen

## Many of Our Stands Are Even-Aged



- Even-aged—trees started at same time after a disturbance
- Some trees in a stand are larger than others—they occupied the site, captured the sun, overtopped others
- Crowns larger, diameter larger—yet trees are same age
- Large diameter trees aren't necessarily older—Diameter not a good predictor of age

## Intermediate Practices

- Tending the crop
- Provide sunlight to the crown
- Young to "middle age" stands
- Improve the existing stand quality
- Provide money, products such as firewood
- Remove insect/diseased trees
- Limited effect structural diversity
- Regeneration not goal- openings too small to encourage germination and sustain seedling/sapling growth

## Intermediate Activities (Tending)

- Release
- Thinning  
(weeding and thinning, crop tree release)
- Improvement  
Cutting
- Pruning



Release

## Thinning

weeding and thinning, tsi or timberstand improvement, fsi or forest stand improvement

- Increase growth of specific trees (crop trees)
- Remove trees deemed less desirable
- 20-50+ years
- 4-10 inches DBH (pole size)
- 10- 16 inches DBH (small sawlog)
- Product firewood



## When thinning What I do when I choose trees to cut


- Look for the trees I want to grow
  - Species – Most valuable for timber- keep options open to cut for timber in the future
  - Healthy – look up at the top, trees with the largest tops relative to their neighbors
  - Relatively straight, with at least one log before trunk forks (maintains option to cut timber in the future)
- Remove trees touching their top
- Mark the trees to cut (or leave)









Leave trees with large, healthy tops.



Leave valuable crop trees: timber, mast, aesthetics





Crop Trees Left to  
Grow


Crop Trees Cut




Leave Healthy Trees

Mark Your Trees to Cut and Leave





## Regeneration (seedling or sprouts)



All about manipulating light

Timing of the harvest is important:

- In terms of the life of the stand- and-
- In terms of time of year
- In good seed years

Which silvicultural technique use depends on:

- species present
- species want
- site capability

## Planting after you cut trees?

In New England-  
not usual

Most of our  
forests  
regenerate  
rapidly after  
cutting



## Sprouts


Most hardwood species  
stump sprout

especially important for  
regenerating red maple and  
red oak



Aspen and Beech  
root sucker/clone

Softwoods do neither and must be  
regenerated from seed



**Remember tree tolerance: Opening size important in determining which species will regenerate.**



**Opening size determines amount of light in the opening**




## Site & Soil Suggest Species to Grow

- white ash, sugar maple → • moderate well drain & enriched fine texture
- beech → • sandy tills
- red oak → • sandy tills & outwash
- white pine → • outwash & sandy tills
- red spruce, hemlock, balsam fir → • shallow pan, poorly drained, outwash, shallow to bedrock

## Two-aged stands

Seed tree cuts, deferred shelterwoods, shelterwood with reserves, clearcuts with reserves can be considered two-aged stands as long as some of the original overstory trees remain in the stand



Enhances vertical and horizontal diversity

Common practice on private land to retain some of the overstory trees indefinitely for aesthetics, wildlife trees, future woody material


## Single Tree Selection

Uneven-aged

- At least 3 distinct age classes free to grow
- Achieved by a series of harvests
- Mature and low quality trees cut in all sizes
- Regenerate tolerant species
- Maintains a mature canopy and vertical structure- a wall of green
- Beech, sugar maple, red spruce, balsam fir, hemlock
- Diameter limit cutting not advised

## Diameter Limit Cutting isn't Selection Harvesting

- Choosing trees to cut based primarily on a minimum diameter- cut larger trees
- Smaller diameter trees aren't necessarily younger
- More likely never got enough sun to grow




## Group Selection


Uneven-aged

- 1/4 to 1/2 acre groups cut
  - Up to 2 acres
- Can think of approaching patch clearcuts
- For regenerating intermediate tolerant species (red oak, white pine, white ash, yellow birch)
- aspen and paper birch (groups approach 1 acre)
- Better scarification

## Group Size Openings



Opening Size (diameter of circle)
1/20 acre----52 foot
1/10 acre---75 foot
1/5 acre---- 105 foot
1/4 acre----117 foot
1/2 acre----166 foot
2/3 acre----200 foot
1 acre-----234 foot







## Shelterwood


Even-aged

- Series of harvests to regenerate
- Harvest removes smaller trees, leaving larger trees to provide correct light conditions and seed source
- Cutting can look light to heavy
- Heavier shade regenerates tolerants (red spruce or hemlock)
- Lower amounts of shade regenerates intermediate tolerants (red oak and white pine)
- Cut overstory when understory regenerated- may be in multiple stages





Sometimes looks like selection- difference is timing of final cut of big trees



Time to remove overwood



## Seed Tree

Even-aged

- Leave 5-10 desirable trees per acre
- For seed, visual relief
- Good source for future snags and super canopy trees
- May leave these for entire rotation



## Clearcut

Even-aged

- Cut everything 2" and greater
- Size depends on objectives and ownership
- Variations- patches and strips
- Regenerates
  - intolerant (paper birch, cherry, aspen/poplar)
  - intermediate (yellow birch and red oak)
  - tolerant with advanced regeneration

