

LOVELL LAKE

2016 SAMPLING HIGHLIGHTS

Station – 2 South

Sanbornville, NH



Blue = Oligotrophic

Yellow = Mesotrophic

Red = Eutrophic

Gray = No Data




Figure 1. Lovell Lake Water Quality (2016)

Station 2 South was used as a reference point to represent the overall Lovell Lake water quality. Water quality data displayed in Tables 1 and 2 are surface water measurements with the exception of the dissolved oxygen data that were collected near the lake bottom.

Table 1. 2016 Lovell Lake Seasonal Averages and NH DES Aquatic Life Nutrient Criteria¹

Parameter	Oligotrophic	Mesotrophic	Eutrophic	Lovell Lake Average (range)	Lovell Lake Classification
Water Clarity (meters)	4.0 – 7.0	2.5 - 4.0	< 2.5	7.5 meters (5.0 – 9.6)	Oligotrophic
Chlorophyll a ¹ (ppb)	< 3.3	> 3.3 – 5.0	> 5.0 – 11.0	2.8 ppb (1.4 – 5.2)	Oligotrophic
Total Phosphorus ¹ (ppb)	< 8.0	> 8.0 – 12.0	> 12.0 – 28.0	6.5 ppb (5.1 – 7.8)	Oligotrophic
Dissolved Oxygen (mg/L)	5.0 – 7.0	2.0 – 5.0	<2.0	1.7 mg/L (0.3 – 5.1)	Eutrophic

* Dissolved oxygen concentrations were measured between 7.5 and 11.0 meters, in the layer of rapidly decreasing temperatures, on August 8, 2016.

Table 2. 2016 Lovell Lake Seasonal Average Accessory Water Quality Measurements

Parameter	Assessment Criteria					Lovell Lake Average (range)	Lovell Lake Classification
	< 10 uncolored	10 – 20 slightly colored	20 – 40 lightly tea colored	40 – 80 tea colored	> 80 highly colored		
Color (color units)	< 10 uncolored	10 – 20 slightly colored	20 – 40 lightly tea colored	40 – 80 tea colored	> 80 highly colored	7.8 color units (5.9 – 9.0)	Uncolored
Alkalinity (mg/L)	< 0.0 acidified	0.1 – 2.0 extremely vulnerable	2.1 – 10 moderately vulnerable	10.1 – 25.0 low vulnerability	> 25.0 not vulnerable	14.9 mg/L (14.2 – 15.6)	Low vulnerability
pH (std units)	< 5.5 suboptimal for successful growth and reproduction		6.5 – 9.0 optimal range for fish growth and reproduction			7.2 standard units (6.7 – 7.6)	Optimal range for fish growth and reproduction
Specific Conductivity (uS/cm)	< 50 uS/cm Characteristic of minimally impacted NH lakes		50-100 uS/cm Lakes with some human influence	> 100 uS/cm Characteristic of lakes experiencing human disturbances		126.4 uS/cm (120.2 – 129.7)	Characteristic of lakes experiencing human disturbances





Figure 2 and 3. Seasonal Secchi disk transparency, chlorophyll a concentrations and dissolved color concentrations. Figures 2 and 3 illustrate the interplay among Secchi Disk transparency, chlorophyll a and dissolved color. Shallower water transparency measurements oftentimes correspond to increases in chlorophyll a and/or color concentrations.

LONG-TERM TRENDS

WATER CLARITY: The Lovell Lake water clarity measurements, measured as Secchi Disk transparency, display a trend of increasing water clarity over the past twenty-eight years (Figure 4).

CHLOROPHYLL: The Lovell Lake chlorophyll *a* concentrations, a measure of microscopic plant life within the lake, display a trend of decreasing concentrations over the past twenty-eight years (Figure 4).

TOTAL PHOSPHORUS: Phosphorus is the nutrient most responsible for microscopic plant growth. The Lovell Lake total phosphorus concentrations display a trend of increasing concentrations over the twenty-eight year span (Figure 5).

COLOR: The Lovell Lake color data, the result of naturally occurring “tea” colored substances from the breakdown of soils and plant materials, have decreased over the past twenty-eight years (Figure 5).

**Figure 4. Lovell Lake - Site 2 South (1989-2016)
Long-term Secchi Disk and Chlorophyll *a* Data**





Table 3. Salmon Falls Headwaters Seasonal Average Water Quality Inter-comparison (2016)

Lake	Average Secchi Disk Transparency (meters)	Average Chlorophyll <i>a</i> (ppb)	Average Total Phosphorus (ppb)	Average Dissolved Oxygen (ppm)
Great East Lake	11.6	1.2	4.4	8.6
Wilson Lake	7.0	2.4	5.5	0.1
Lovell Lake	7.5	2.8	6.5	1.7
Horn Pond	7.3	1.9	5.7	2.9
Lake Ivanhoe	4.1	3.9	7.9	-----

- Water quality data are reported for a deep reference sampling location in each water body
- Dissolved oxygen measurements were collected in the summer (late July and August) in the bottom water layer (metalimnion or hypolimnion).
- ----- Indicates the site is too shallow to form a bottom water layer (metalimnion or hypolimnion) during the summer months.


**Figure 5. Lovell Lake - Site 2 South (1989-2016)
Long-term Total Phosphorus and Dissolved Color Data**



Figures 4 and 5. Changes in the Lovell Lake water clarity (Secchi Disk depth), chlorophyll *a*, dissolved color and total phosphorus concentrations measured between 1989 and 2016. **These data illustrate the relationship among plant growth, dissolved color and water clarity. Total phosphorus data are also displayed and are oftentimes correlated with the amount of plant growth.**

Figure 6. Monthly Lovell Lake dissolved oxygen profiles collected between May 23 and October 3, 2016. The vertical red line indicates the oxygen concentration commonly considered the threshold for successful growth and reproduction of cold water fish such as trout and salmon. *Notice the low oxygen concentrations near the lake bottom between July 25 and September 19.*

**Figure 6. Lovell Lake, Site 2 South - Dissolved Oxygen Profiles
(May 23 through October 3, 2016)**



Recommendations


Implement Best Management Practices within the Lovell Lake watershed to minimize the adverse impacts of polluted runoff and erosion on the lake. Refer to “Landscaping at the Water’s Edge: An Ecological Approach” and “New Hampshire Homeowner’s Guide to Stormwater Management: Do-It-Yourself Stormwater Solutions for Your Home” for more information on how to reduce nutrient loading caused by overland run-off. The Acton Wakefield Watershed Alliance also offers technical assistance to help design and implement erosion control projects that protect and improve the water quality.

- http://extension.unh.edu/resources/files/Resource004159_Rep5940.pdf
- <http://soaknh.org/wp-content/uploads/2016/04/NH-Homeowner-Guide-2016.pdf>
- <http://awwatersheds.org/healthy-lakes/conservation-practices-for-homeowners/>

Figure 7. Lovell Lake

Sanbornville, NH

2016 Deep water sampling site locations that display the seasonal average water clarity



0 0.2 0.4 0.6 0.8 Miles



Extension



Aerial Orthophoto Source: NH GRANIT
Site location GPS coordinates collected by the UNH Center for Freshwater Biology