

Pine-Oak-Hemlock in the Northeast

Silvics, Dynamics, Climate Change, Restoration

Tom Lee

Department of Natural Resources and the Environment University of New Hampshire



Topics


- Where is pine-oak-hemlock forest?
- Silvics of individual species
- Succession and stand dynamics, including the effects of land use history
- Climate change adaptation and restoration


Forest Types of New England


(Westveld et al. 1956)


"Transition hardwoods – white pine – hemlock"


"Central hardwoods – hemlock – white pine"

































- Where is pine-oak-hemlock forest?
- Silvics of individual species
 - Site affinity
 - Life history characteristics
- Succession and stand dynamics, including the effects of land use history
- Climate change adaptation and restoration

Eastern White Pine


- Naturally occurs on excessively drained soils
 - Coarse texture (sands, sandy loams)
 - Glacial outwash
 - Dry, nutrient poor
- Pine is drought tolerant
 - Not as much as oaks
 - Produces "sinker roots" in deep coarse soils (up to 15' deep)



Leak and Riddle (1979)

Eastern White Pine

- Occurs differentially on poorer soils – BUT – growth is better on mesic, fertile soils
- Limited to poorer soils by vigorous competition elsewhere



Eastern Hemlock

- Associated with shallow soils
 - Shallow to bedrock or hard pan
 - Rocky, often just loose rock
- Soils can be wet
- Sometimes on sandy soils
- BUT: as in pine, growth is better on mesic, fertile soils







Northern Red Oak

- In southern, central parts of region
 - Often on middle and lower slopes
 - Soils typically derived from fine, rocky till
 - Soils from clay to loamy sands


Competition from other mesic hardwoods






Northern Red Oak: Interactions

- Grows best on moist enriched sites
 - But competitively displaced by sugar maple
- On moist, fertile till sites: Displaces pine
- Dry, less fertile sites: Outgrown by pine






Black Oak, White Oak





Also occur on dry, coarse-textured soils



- More tolerant of stress than red oak
- Displaced to warmer, drier, less fertile sites by COMPETITION with red oak and others
- Mix with white pine on coarse soils

Red Maple

- Incredibly broad tolerances to moisture level and soil fertility
- Persists in standing water
 - Forms pure stands in flooded areas
- Occurs with black and white oak on ridges





Sweet Birch

- Avoids extremely dry or wet sites
- Grows on shallow rocky soils, deep sandy soils





American Beech

- Associated with deep, well-drained coarse or washed glacial till
- Not as flood tolerant as red maple or ash
- Reduced on enriched sites by competition with sugar maple






Leak and Riddle (1979)

Sugar Maple, White Ash

- Associated with moist, fertile ("enriched") sites
 - Less acidic
 - High levels of calcium
- Lower slopes, coves, fine marine deposits



Leak and Riddle (1979)


Paper Birch, Gray Birch, Quaking (& bigtooth) Aspen


• Fairly broad tolerances to physical factors







- Where is pine-oak-hemlock forest?
- Silvics of individual species
 - Site affinity
 - Life history characteristics
- Succession and stand dynamics, including the effects of land use history
- Climate change adaptation and restoration









Regeneration characteristics

- ROOT SPROUTERS
 - □ Aspen
- (+) USEFUL IN PROMOTING ASPEN AFTER LOGGING
- Beech
- (-) MAY DOMINATE UNDERSTORY
- (-) ALLOWS POST-LOGGING RESPONSE
- STUMP SPROUTERS
 - Most of the hardwoods:
 OAKS, RED MAPLE, SWEET BIRCH

Regeneration characteristics

• STUMP SPROUTERS





Eli Sagor, University of MN Extension

Growth characteristics

- Eastern white pine
 - Height growth slow first few years

Third year






Fifth year

Growth characteristics

- Northern red oak
 - Remarkable height and radial growth
 - Can overtop other mid-tolerant species of same age







50-year radial 107 75 68 growth (mm)


- Where is pine-oak-hemlock forest?
- Silvics of individual species
- Succession and stand dynamics, including the effects of land use history
- Climate change adaptation and restoration

Stand dynamics (focus on uplands)

- Old-field succession
- Succession after clear-cutting
- Effects of smaller openings on stand dynamics















Why pine successful in old fields?


- 1) Pine competes well with herbaceous plants
- 2) Took time for other woody plants to arrive on-site and compete
- 3) Meadow voles a common herbivore prefer hardwood seedlings



































Relevance to silvicultural systems

- Large openings (> 1/4 acre)
 - Regenerate most species
 - Required for paper birch, aspen
 - Allow faster growth of oak and pine



Relevance to silvicultural systems

- Group selection
 - Favors mid-tolerant hardwoods
 - Especially sweet birch, red maple, red oak
 - To some extent, pine (depends on site, size)





Relevance to silvicultural systems


- Single-tree selection
 - Favors shade tolerant species (beech, sugar maple, hemlock)









- Where is pine-oak-hemlock forest?
- Silvics of individual species
- Succession and stand dynamics, including the effects of land use history
- Climate change adaptation and restoration








Climate change: Concerns

- OPERATIONS
 - Shorter duration of frozen ground
 - Rainfall events impact infrastructure, access
- DISTURBANCE (lots of uncertainty here)
 - Increased storm intensity = more flooding?
 - Drought = more frequent fire?
 - More frequent impact on stand structure?
- DIRECT EFFECTS ON TREES
 - Productivity may increase
 - Tree species' DISTRIBUTIONS and ABUNDANCES



USFS Climate Change Tree Atlas

"...we are predicting potential suitable habitat by year 2100 ... NOT ... where the species will be at that time, as great lag times are involved ...

...the model does not account for future biotic interactions (competition...) or other human ... or natural ... disturbances ..."

Pine-oak-hemlock region: Projected changes in abundance

DECREASE INCREASE LITTLE CHANGE
Eastern white pine Black oak Sweet birch

Northern red oak White oak

American beech Scarlet oak DEPENDS ON MODEL Hickories Red maple

Sugar maple Yellow poplar White ash

Paper birch Aspen

DECREASE ≠ DISAPPEARANCE! TRENDS?

Pine-oak-hemlock to oak-hickory but at the same time

Nor. hardwoods to pine-oak-hemlock

Changes in tree composition will likely occur *slowly* "GREAT LAG TIMES"

- Trees have broad responses to temperature
 - predicted temperature changes will not kill trees outright
- Trees are long-lived; hold space
- Migration of trees across landscape is slow
- UNKNOWNS: Disturbance, insects, and disease
 - Hemlock woolly adelgid and temperature

NO REASON TO MAKE HUGE CHANGES IN HOW WE MANAGE OUR FORESTS

How can we minimize negative effects of climate change on forestry in the pine-oak-hemlock region?

"CLIMATE CHANGE ADAPTATION"







Three ADAPTIVE approaches

(Millar et al. 2007)


- **RESISTANCE** options
 - RESISTANCE: Ability of a forest to remain the same in the face of some environmental change or disturbance
- RESILIENCE options
 - RESILIENCE: Ability of a forest stand to recover from impact and return to its previous state
- **RESPONSE** options
 - RESPONSE: transition of forest stand from present state to new, climate-adjusted state

Consider two scales:

INDIVIDUAL STAND



LANDSCAPE



Gunn et al. (2009)

STAND LEVEL Climate change RESISTANCE

- How to make a stand more RESISTANT to climate change?
- *Reduce* OTHER stresses
 - Be pro-active in minimizing insect pest impacts
 - MONITORING!
 - Appropriate thinning to reduce competition, improve individual tree health





LANDSCAPE LEVEL Climate change RESISTANCE

- How to make a stands in a landscape more RESISTANT?
- Reduce OTHER stresses
 - Maintain mature stands as wind protection for managed stands

STAND LEVEL Climate change RESILIENCE

- Best predictor of RESILIENCE is diversity
- A more diverse stand is more resilient to climate change

WHY?

STAND LEVEL Climate change RESILIENCE

PORTFOLIO EFFECT (from finance)

- A greater DIVERSITY of investments means you are more likely to survive changes in the economy
- While some investments do *poorly* in periods of change, others will do *well*



STAND LEVEL Climate change RESILIENCE

A stand with

high number of species or high diversity of age classes, or with species having high gene diversity

Should be more **RESILIENT**



Consistent with mixedstand approaches (Kelty et al.)



LANDSCAPE LEVEL Climate change RESILIENCE

A landscape with

high diversity of forest types
high diversity of stand ages, or
high diversity of stand structures
is likely more **RESILIENT** to climate change

Climate change RESILIENCE

SO, FOR RESILIENCE: Important to maintain biotic, age, and structural diversity at all scales



Three ADAPTIVE approaches

(Millar et al. 2007)

- **RESISTANCE** options
 - RESISTANCE: Ability of a forest to remain the same in the face of some environmental change or disturbance
- **RESILIENCE** options
 - RESILIENCE: Ability of a forest stand to recover from impact and return to its previous state
- **RESPONSE** options
 - RESPONSE: transition of forest stand from present state to new, climate-adjusted state

Climate change RESPONSE

Here, we FACILITATE CHANGE TO NEW FOREST COMMUNITY


- LANDSCAPE LEVEL
 - Promote connectivity of stands
 - Promote diversity of communities
- STAND LEVEL
 - Shorten rotation times

Topics

- Where is pine-oak-hemlock forest?
- Silvics of individual species
- Succession and stand dynamics, including the effects of land use history
- Climate change adaptation and restoration









What can we do about invasive species?

- Prevention
 - Effective, involves politics and policy
- Early detection, rapid response (EDRR)
 - Quickly identify new invasive species
 - Find populations and reduce them
 - Requires MONITORING!







Eradication Control

- Herbicides
- Mechanical approaches





Topics

- Where is pine-oak-hemlock forest?
- Silvics of individual species
- Succession and stand dynamics, including the effects of land use history
- Climate change adaptation and restoration



