

NATICOOK LAKE

2019 SAMPLING HIGHLIGHTS

Station – 1 Deep

Merrimack, NH



Extension

Blue = Oligotrophic

Yellow = Mesotrophic

Red = Eutrophic

Gray = Not Assessed




Figure 1. Naticook Lake Water Quality (2018)

Station 1 Deep (Figure 8) was used as a reference point to represent the overall Naticook Lake water quality. Water quality data displayed in Tables 1 and 2 are surface water measurements with the exception of the Dissolved Oxygen data that were collected towards the lake bottom.

Table 1. 2019 Naticook Lake Seasonal Averages and NH DES Aquatic Life Nutrient Criteria¹

Parameter	Oligotrophic	Mesotrophic	Eutrophic	Naticook Lake Average (range)	Naticook Lake Classification
Water Clarity (meters)	4.0 – 7.0	2.5 - 4.0	< 2.5	4.1 meters (2.9 – 5.1)	Oligotrophic
Chlorophyll a ¹ (ppb)	< 3.3	> 3.3 – 5.0	> 5.0 – 11.0	1.8 ppb (1.3 – 2.6)	Oligotrophic
Total Phosphorus ¹ (ppb)	< 8.0	> 8.0 – 12.0	> 12.0 – 28.0	12.0 ppb (11.4 – 12.6)	Mesotrophic
Dissolved Oxygen (mg/L)	5.0 – 7.0	2.0 – 5.0	<2.0	0.7 mg/L (0.0 – 2.9) *	Eutrophic *

* Dissolved oxygen concentrations were measured on August 30, between 3.5 and 5.5 meters, in the layer of rapidly decreasing temperature.

Table 2. 2019 Naticook Lake Seasonal Average Accessory Water Quality Measurements

Parameter	Assessment Criteria					Naticook Lake Average (range)	Naticook Lake Classification
Color (color units)	< 10 uncolored	10 – 20 slightly colored	20 – 40 lightly tea colored	40 – 80 tea colored	> 80 highly colored	32.2 color units (range: 28.3 – 34.7)	Lightly tea colored
Alkalinity (mg/L)	< 0.0 acidified	0.1 – 2.0 extremely vulnerable	2.1 – 10 moderately vulnerable	10.1 – 25.0 low vulnerability	> 25.0 not vulnerable	19.6 mg/L (range: 18.8 – 20.2)	Low Vulnerability
pH (std units)	< 5.5 suboptimal for successful growth and reproduction		6.5 – 9.0 optimal range for fish growth and reproduction			7.1 standard units (single value)	Optimal range for fish growth and reproduction
Specific Conductivity (uS/cm)	< 50 uS/cm Characteristic of minimally impacted NH lakes		50-100 uS/cm Lakes with some human influence	> 100 uS/cm Characteristic of lakes experiencing human disturbances		257.6 uS/cm (single value)	Characteristic of lakes experiencing human disturbances





Figure 2 and 3. Seasonal Secchi Disk transparency, chlorophyll a changes, and dissolved color concentrations. Figures 2 and 3 illustrate the interplay among Secchi Disk transparency, chlorophyll a, and dissolved color. Shallower water transparency measurements oftentimes correspond to increases in chlorophyll a and/or color concentrations.


LONG-TERM TRENDS

WATER CLARITY: The Naticook Lake water clarity measurements, measured as Secchi Disk transparency, have oscillated among years while the long-term water clarity trend is stable (Figure 4).

CHLOROPHYLL: The Naticook Lake chlorophyll *a* concentrations, a measure of microscopic plant life within the lake, have oscillated among years while the long-term trend is stable (Figure 4).

TOTAL PHOSPHORUS: Phosphorus is the nutrient most responsible for microscopic plant growth. The Naticook Lake total phosphorus concentrations display a trend of increasing concentrations (Figure 5).

COLOR: The Naticook Lake color data, the result of naturally occurring “tea” color substances from the breakdown of soils and plant materials, display a trend of increasing concentrations (Figure 5).



Figures 4 and 5. Changes in the Naticook Lake water clarity (Secchi Disk depth), chlorophyll *a*, dissolved color, and total phosphorus concentrations measured between 1983 and 2019. **These data illustrate the relationship among plant growth, water color, and water clarity. Total phosphorus data are also displayed and are oftentimes correlated with the amount of plant growth.** Long-term trends are based on the analysis of annual median values.






Figure 6. Naticook Lake surface water total phosphorus inter-site comparison. *Notice the difference in total phosphorus concentrations among sampling locations.*

Figure 7. Naticook Lake dissolved oxygen profile collected on August 30, 2019. The vertical red line indicates the dissolved oxygen concentration commonly considered the threshold for successful growth and reproduction of warm water fish such as bass and perch. *Notice the low dissolved oxygen concentrations near the lakebottom.*



Recommendations

Implement Best Management Practices within the Naticook Lake watershed to minimize the adverse impacts of polluted runoff and erosion on Naticook Lake. Refer to “Landscaping at the Water’s Edge: An Ecological Approach” and “New Hampshire Homeowner’s Guide to Stormwater Management: Do-It-Yourself Stormwater Solutions for Your Home” for more information on how to reduce nutrient loading caused by overland run-off.



- https://extension.unh.edu/resources/files/Resource004159_Rep5940.pdf
- <https://www.des.nh.gov/organization/commissioner/pip/publications/wd/documents/wd-11-11.pdf>

Figure 8. Naticook Lake
Merrimack, NH
2018 Deep and near-shore sampling sites.

Average Depth = 5.9 feet
Maximum Depth = 19.1 feet
Surface Area = 62.4 acres

Disclaimer: Due to the COVID-19 pandemic, access to University files needed to update this image, the 2018 map was used. The sampling locations, average depth, maximum depth and surface area are accurately depicted. However, the year and seasonal average water transparency have not been updated. Once full University resources become available this map will be updated.

The information in the first two pages of this report reflects the 2019 conditions and is final.



Aerial Orthophoto Source: NH GRANIT
Site location GPS coordinates collected by the UNH Center for Freshwater Biology

