MARSH POND

2019 SAMPLING HIGHLIGHTS

Station – 2 Deep

New Durham, NH



Station 2 Deep (Figure 5) was used as a reference point to represent the overall Marsh Pond water quality. With the exception of the dissolved oxygen measurements, the water quality data displayed in Tables 1, 2 and 3 are surface water measurements.




Figure 1. Marsh Pond Water Quality (2019)

Table 1. 2019 Marsh Pond Seasonal Averages and NH DES Aquatic Life Nutrient Criteria¹

Parameter	Oligotrophic	Mesotrophic	Eutrophic	Marsh Pond Average (range)	Marsh Pond Classification
Water Clarity (meters)	4.0 - 7.0	2.5 - 4.0	< 2.5	2.8 meters (2.1 – 3.6)	Mesotrophic
Chlorophyll a ¹ (ug/l)	< 3.3	> 3.3 - 5.0	> 5.0 - 11.0	14.2 <i>u</i> g/l (4.8 – 24.6)	Eutrophic
Total Phosphorus ¹ (ug/l)	< 8.0	> 8.0 - 12.0	> 12.0 - 28.0	48.4 <i>u</i> g/l (19.8 – 80.8)	Eutrophic
Dissolved Oxygen (mg/L)	5.0 - 7.0	2.0 - 5.0	<2.0	* 2.0 mg/l (0.0 – 4.7)	* Mesotrophic

* Marsh Pond developed a mid-lake layer of rapidly decreasing temperature and the July 24 oxygen profile was used to classify Marsh Pond.

Table 2. 2019 Marsh Pond Seasonal Average Accessory Water Quality Measurements

Parameter	Assessment Criteria					Marsh Pond Average (range)	Marsh Pond Classification
Color (color units)	< 10 uncolored	10 – 20 slightly colored	20 – 40 lightly tea colored	40 – 80 tea colored	> 80 highly colored	19.3 color units (range: 12.7 – 22.2)	Slightly tea colored
Alkalinity (mg/L)	< 0.0 acidified	0.1 – 2.0 extremely vulnerable	2.1 – 10 moderately vulnerable	10.1 – 25.0 Iow vulnerability	> 25.0 not vulnerable	7.9 mg/L (range: 7.1 – 8.7)	Moderately vulnerable
pH (std units)	suboptimal	5.5 for successful reproduction	6.5 – 9.0 optimal range for fish growth and reproduction		6.9 standard units (range: 6.9 – 7.0)	Optimal range for fish growth and reproduction	
Specific Conductivity (<i>u</i> S/cm)	Characteristi	<i>u</i> S/cm c of minimally d NH lakes	50-100 <i>u</i> S/cm Lakes with some human influence	> 100 <i>u</i> Characterist experiencir disturba	ic of lakes ng human	53.8 <i>u</i> S/cm (range: 48.3 – 56.9)	Characteristic of lakes with some human influence




Figure 2 and 3. Seasonal Secchi disk transparency, chlorophyll a concentrations and dissolved color concentrations. Figures 2 and 3 illustrate the interplay among Secchi Disk transparency, chlorophyll a and dissolved color. Shallower water transparency measurements oftentimes correspond to increases in chlorophyll a and/or color concentrations.

Lake	Average (range) Secchi Disk Transparency (meters)	Average (range) Chlorophyll <i>a</i> (ppb)	Average (range) Total Phosphorus (ppb)	Average (range) Dissolved Color (CPU)	Average (range) Dissolved Oxygen (mg/l)
Merrymeeting Lake	10.2 meters	0.7 <i>u</i> g/l	3.7 <i>u</i> g/l	4.2 CPU	10.9 mg/l
	(range: 8.2 – 12.5)	(range: 0.1 – 1.0)	(range: 2.6 – 6.5)	(range: 1.4 – 7.4)	(range: 8.7 – 12.2)
Marsh Pond	2.8 meters	14.2 ug/l	48.4 <i>u</i> g/l	19.3 CPU	2.0 mg/l
	(range: 2.1 – 3.6)	(range: 4.8 – 24.6)	(range: 19.8 – 80.8)	(range: 12.7 – 22.2)	(range: 0.0 – 4.7)
Jones Pond	2.7 meters	10.5 ug/l	29.0 <i>u</i> g/l	29.6 CPU	0.2 mg/l
	(range: 2.1 – 3.5)	(range: 4.0 – 13.8)	(range: 18.2 – 34.8)	(range: 24.3 – 35.6)	(range: 0.0 – 0.3)
Downing Pond	2.7 meters (range: 2.2 – 3.0)	8.0 <i>u</i> g/l (range: 4.1 – 12.2)	28.7 <i>u</i> g/l (range: 19.0 – 37.4)	35.2 CPU (range: 25.7 – 41.1)	


• Water quality data are reported for a deep reference sampling location in each lake/pond.

• Dissolved oxygen measurements were collected in the summer (mid to late July) in the bottom water layer (hypolimnion or metalimnion).

Downing Pond Secchi Disk transparency measurements intermittently reached the lake bottom before disappearing from view and likely underestimate the water transparency.

------ Indicates the site is too shallow to form a stable deep water layer (hypolimnion or metalimnion) during the summer months.

Figure 4. Marsh Pond dissolved oxygen profiles collected between May 19 and September 4, 2019. The vertical red line indicates the oxygen concentration commonly considered the threshold for successful growth and reproduction of cold water fish such as brook trout. *Note: low dissolved oxygen concentrations near the lake bottom generally corresponded to elevated deep water total phosphorus concentrations.*




Recommendations

Review the "Merrymeeting Lake & River Watershed Management Plan" that provides background information and offers potential solutions to existing water quality problems. Homeowners within the Marsh Pond watershed should consider implementing Best Management Practices to minimize the adverse impacts of polluted runoff and erosion on Marsh Pond. Homeowners can refer to "Landscaping at the Water's Edge: An Ecological Approach" and "New Hampshire Homeowner's Guide to Stormwater Management: Do-It-Yourself Stormwater Solutions for Your Home". Both self-help documents offer relatively simple solutions to reduce nutrient loading caused by overland run-off.

- <u>https://www.newdurhamnh.us/home/news/merrymeeting-lake-river-watershed-management-plan</u>
- <u>https://extension.unh.edu/resources/files/Resource004159_Rep5940.pdf</u>
- https://www.des.nh.gov/organization/commissioner/pip/publications/wd/documents/wd-11-11.pdf

Figure 5. Marsh Pond

New Durham, NH 2018 Deep water sampling site



Aerial Orthophoto Source: NH GRANIT GPS Coordinates collected by the UNH Center for Freshwater Biology



