



Raising Broilers



Broilers are hybrid chickens bred for fast growth and finish. Typically raised to about 5 pounds, they are usually processed between 7 and 9 weeks of age. Birds slaughtered between 9 and 12 weeks of age, typically weigh between 7 and 10 lbs. and are called roasters.

It's possible to enjoy a continuous supply of chicken by starting a new flock of broiler chicks when the previous flock is a month old. This way you can process one-quarter of each flock at seven, eight, nine and 10 weeks of age. How many pounds of chicken do you eat a week? This is a good way to determine how many to raise a year. Remember that one 21 cubic foot freezer holds approximately 850 pounds of meat.

Housing

Start chicks indoors in a deep container (at least 30 inches deep) which will supply one square foot of floor space per bird (plastic Sterlite containers work well, but if the container is square, round the corners in some way to prevent chick piling and death). The chicks will require one square foot of floor space for approximately 1 month at which time the space should double.

Spread an absorbent litter on the floor of the box that is not small enough to be mistaken for food. If newspaper is used as bedding, sprinkled it with a **thin** layer of sterile sand or leg splaying can occur.

Replace wet soiled bedding/litter as often as needed to keep chicks clean and dry.

UNH Cooperative Extension Programs

	Community and Economic Development
	Food and Agriculture ✓
	Natural Resources
	Youth and Family

Broilers are hybrid chickens bred for fast growth and finish.



Photo captions are Myriad Pro, 9pt, Bold

Brooding

Chicks cannot self-regulate their body temperature until they are 14 days old. This means that in their first two weeks of life they are extremely sensitive to variations in temperature and that they are easily overheated or chilled. Maintain a temperature at chick level of approximately 90 to 95 degrees F. Don't guess! Use a thermometer. An infrared bulb securely suspended from above that can be lowered or raised as needed to regulate the ambient air temperature should be used. Incandescent lights can also be used. **REMEMBER these heat sources combined with the dry bedding can cause fire so take proper precautions to keep your family and animals safe.** Drop the temperature 5 degrees per week until you reach 70 degrees at which time birds should be fledged so the lights should no longer be needed at night (approximately 5 weeks of age depending on the season of the year and facilities birds are kept in. Contact your local extension for more information).

The most important nutrient to provide new chicks with is water.

Feeding and Watering

The most important nutrient to provide new chicks with is water. Just before hatching, chicks absorb the rest of the egg yolk contained in their egg. This nutrient rich food will sustain them for approximately 3 days. Therefore, making sure water is in place in the brooder for when the chicks arrive is more important than having food available. As each bird goes into the brooder, dip the beak into the water so they can get a small drink and hopefully learn where the water source is. Placing colored glass marbles in the trough can encourage chicks to peck at the water and learn to drink from it. Monitor the birds carefully at first and when you believe all have learned to drink, then you can put the food in the brooder (no later than the end of the 2nd day). These steps can prevent chicks from getting pasting by preventing the birds from filling up on feed and not drinking the water like they should. **Without water in their system, they cannot correctly digest their food, which leads to pasting. Incorrect brooder temperatures is also another factor that can lead to pasting.**

To be successful raising broilers, use medicated chick starter when you get the chicks home (100lbs of starter is enough to start 25 broiler chicks). After 2 weeks birds can go on broiler grower and at no less than 42 days birds can go on finisher.

Open containers like cardboard egg cartons can be used to supply food for the first three or four days but these open containers should be switched to limited access containers to prevent chicks from standing in feed and defecating in it. Proper chick feeders are readily available at feed stores.

Food and water should be available to broilers constantly until slaughter because of the rapid growth rate these birds experience. Even pastured birds should be supplied with free choice rations and water. Pasture should only be used to supplement a balanced ration/ pelleted feed.

A final note:

Consult your local zoning and building ordinances before beginning any household livestock operation. Laws and ordinances in some communities (even rural communities) may prohibit or restrict such activities in your neighborhood, regardless of land mass. One should also consider the impact of your home livestock operation on your neighbors. Use care in location and construction of housing for your animals and develop a plan for manure management that will prevent odor, environmental problems, and adhere to NH Department of Agriculture, Markets, and Food, Best Management Practices.

Created: August 2000
Updated: January 2018

Visit our website:
extension.unh.edu

UNH Cooperative Extension brings information and education into the communities of the Granite State to help make New Hampshire's individuals, businesses, and communities more successful and its natural resources healthy and productive. For 100 years, our specialists have been tailoring contemporary, practical education to regional needs, helping create a well-informed citizenry while strengthening key economic sectors.

The University of New Hampshire Cooperative Extension is an equal opportunity educator and employer. University of New Hampshire, U.S. Department of Agriculture and N.H. counties cooperating.

About the Authors

Original fact sheet written by Tom Danko, retired Extension Poultry Specialist. Revised October 17, 2017 by Dot Perkins, UNHCE Field Specialist-Dairy, Forages and Livestock Team.

For More Information

State Office

Taylor Hall
59 College Rd.
Durham, NH 03824
<http://extension.unh.edu>

Education Center and Infoline

answers@unh.edu
1-877-EXT-GROW
(1-877-398-4769)
9 a.m. to 2 p.m. M-F
extension.unh.edu/askunhex-tension